

Product Information

Version 2.0


Voltaren Emulgel diclofenac diethylammonium 1.16% w/w gel AUST R 47676

Voltaren Osteo Gel diclofenac diethylammonium 1.16% w/w gel AUST R 175889

Voltaren Emulgel & Voltaren Osteo Gel

Name of the medicine

Active ingredient: diclofenac diethylammonium 11.6 mg/g



Chemical structure:

CAS Registry number: 78213-16-8

Description

Voltaren Emulgel/ Voltaren Osteo Gel contains diethylammonium-[o-((2,6-dichlorophenyl)-amino)-phenyl]-acetate) (=diclofenac diethylammonium),

One gram of the gel contains 11.6 mg diclofenac diethylammonium equivalent to 10 mg diclofenac sodium.

Voltaren Emulgel / Voltaren Osteo Gel contains the excipients: carbomer 934P, cetomacrogol 1000, coco-caprylate/caprate, diethylamine, isopropyl alcohol, paraffin – liquid, perfume (containing benzyl benzoate), propylene glycol, water - purified.

Pharmacology

Pharmacodynamics

Pharmacotherapeutic group: Topical products for joint and muscular pain, anti-inflammatory preparations, non-steroids for topical use (ATC code M02A A15).

Diclofenac is a non-steroidal anti-inflammatory drug (NSAID) with pronounced analgesic, anti-inflammatory and antipyretic properties. Inhibition of prostaglandin synthesis is the primary mechanism of action of diclofenac.

Voltaren Emulgel / Voltaren Osteo Gel is an anti-inflammatory and analgesic preparation designed for external application.

In inflammation and pain of traumatic or rheumatic origin, Voltaren Emulgel and Voltarne Osteo Gel has been shown to relieve pain, reduce oedema, and shorten the time to return of normal function.

Due to an aqueous-alcoholic base, the gel also exerts a soothing and cooling effect.

Pharmacokinetics

Absorption:

When Voltaren Emulgel/Voltaren Osteo Gel is applied locally, the active substance is absorbed through the skin. The amount of diclofenac absorbed through intact skin is proportional to the contact time and

skin area covered with the gel, and depends on the total topical dose and the hydration of the skin. Absorption amounts to about 6% of the dose of diclofenac after topical application of 2.5 g the gel per 500cm² skin, determined by reference to the total renal elimination compared with Voltaren tablets. Occlusion over a period of 10 hours leads to a three-fold increase in the amount of diclofenac absorbed. No information is available on the clinical effects and consequences of use under occlusion.

Distribution:

After topical administration of Voltaren Emulgel/Voltaren Osteo Gel to hand and knee joints, diclofenac can be measured in plasma, synovial tissue, and synovial fluid. Maximum plasma concentrations of diclofenac after topical administration of Voltaren Emulgel/Voltaren Osteo Gel is approximately 100 times lower than after oral administration of the same quantity of diclofenac. Diclofenac is highly bound to serum proteins (99.7%), predominantly to albumin (99.4%).

Metabolism:

The biotransformation of diclofenac involves partly glucuronidation of the intact molecule, but mainly by single and multiple hydroxylation and methoxylation, resulting in several phenolic metabolites, most of which are converted to glucuronide conjugates. Two of these phenolic metabolites are biologically active, however, to a much smaller extent than diclofenac. Metabolism of diclofenac following topical and oral administration is similar.

Elimination:

The total systemic clearance of diclofenac from plasma is 263 ± 56 mL/min (mean value ± SD). The terminal plasma half-life is 1 to 2 hours. Four of the metabolites, including the two active ones, also have short plasma half-lives of 1-3 hours. One metabolite, 3'-hydroxy-4'-methoxy-diclofenac, has a much longer plasma half-life. However, this metabolite is virtually inactive. Diclofenac and its metabolites are excreted mainly in the urine.

Special population:

Patients with renal impairment

No accumulation of diclofenac and its metabolites is to be expected in patients suffering from renal impairment.

Patients with hepatic impairment

In patients with chronic hepatitis or non-decompensated cirrhosis, the kinetics and metabolism of diclofenac are the same as in patients without liver disease.

Preclinical safety data

Preclinical data from acute and repeated dose toxicity studies, as well as from genotoxicity, mutagenicity, and carcinogenicity studies with diclofenac revealed no specific hazard for humans at the intended therapeutic doses. The gel was well tolerated in a variety of studies. There was no potential for phototoxicity and Voltaren Emulgel/Voltaren Osteo Gel caused no skin sensitisation.

Indications

For the short term (up to 2 weeks) local symptomatic treatment of the following musculoskeletal inflammatory conditions:

- acute soft-tissue injuries, including sprains, strains and sports injuries
- localised forms of soft tissue rheumatism such as tendinitis (eg tennis elbow) and bursitis.

For the short term (up to 3 weeks) relief of pain in non-serious arthritis (i.e. mild and localised forms of osteoarthritis) of the knees or fingers. Relief of osteoarthritic pain builds up gradually over the first few days of treatment; a significant effect can be expected after one week of application.

Contraindications

- Known hypersensitivity to diclofenac or to any of the other ingredients in the gel.
- Patients in whom attacks of asthma, urticaria, or acute rhinitis are precipitated by aspirin or other non-steroidal anti-inflammatory drugs (NSAIDs) (see 'Precautions – Pre-existing asthma').
- Last trimester of pregnancy (see 'Precautions – Use in pregnancy').

Precautions

Systemic side effects

The likelihood of systemic side effects occurring following topical diclofenac is small compared with the frequency of side effects following oral diclofenac. However, the possibility of systemic side effects cannot be excluded, particularly when the gel is applied to relatively large areas of skin or for periods longer than 3 weeks. In case such usage is envisaged, the product information on Voltaren Rapid should be consulted.

Topical side effects

Voltaren Emulgel/Voltaren Osteo Gel should be applied only to intact, healthy skin and not to skin wounds, infections, exudative dermatoses or open injuries. It should not be allowed to come into contact with the eyes or mucous membranes.

Voltaren Emulgel/Voltaren Osteo Gel contains propylene glycol and benzyl benzoate, which may cause mild, localised skin reactions in some people.

Use in gastrointestinal disorders

In general, topical NSAIDs should be used with caution in those patients with a history of (or active) gastro-intestinal ulceration or bleeding,

Use in renal impairment

In general, topical NSAIDs should be used with caution in those patients with a history of severe renal impairment.

Use in pre-existing asthma

Reactions to NSAIDs such as asthma exacerbations (so-called intolerance to analgesics / analgesics-asthma), Quincke's oedema or urticaria are more frequent in patients with asthma, seasonal allergic rhinitis, swelling of the nasal mucosa (i.e. nasal polyps), chronic obstructive pulmonary diseases or chronic infections of the respiratory tract (especially if linked to allergic rhinitis-like symptoms) than in other patients. Therefore, special precaution is recommended in such patients. This

is applicable as well for patients who are allergic to other substances, e.g. with skin reactions, pruritus or urticaria.

Use under dressings

Voltaren Emulgel/Voltaren Osteo Gel can be used with non-occlusive bandages but should not be used with occlusive dressing (see 'Absorption').

Use in pregnancy (Category C)

The use of diclofenac in pregnant women has not been studied; therefore, Voltaren Emulgel should not be used during the first two trimesters of pregnancy or in women who are likely to become pregnant, unless the potential benefit to the mother outweighs the risk to the foetus.

Diclofenac is contraindicated during the third trimester of pregnancy, owing to the possibility of uterine inertia and/or premature closure of the ductus arteriosus.

Animal studies have not shown any direct or indirect harmful effects on pregnancy, embryonal/foetal development, parturition or postnatal development.

Animal studies did see some degree of reproductive toxicity, although this was generally associated with maternal toxicity, with no indication of developmental toxicity.

Genotoxicity and carcinogenicity

There was no evidence that diclofenac had a teratogenic potential in mice, rats or rabbits.

Effects on fertility

Diclofenac had no influence on the fertility of parent animals in rats. The prenatal, perinatal and postnatal development of the offspring was not affected.

Use in lactation

It is not known whether topical diclofenac is excreted in breast milk; therefore, Voltaren Emulgel is not recommended during breastfeeding. If there are compelling reasons for using it, it should not be applied to the breasts or to large areas of skin, nor should it be used for a prolonged period.

Use in children

Voltaren Emulgel/Voltaren Osteo Gel is not recommended for use in children under 12 years of age, as safety and efficacy in this age group have not been established.

Use in the elderly

No dose adjustment is required for elderly patients.

Effects on ability to drive and use machines

Cutaneous application of the gel is unlikely to influence on the ability to drive and use machines.

Interactions with other medicines

Anticoagulants and antiplatelet agents:

The concurrent use of NSAIDs and warfarin has been associated with severe, sometimes fatal, haemorrhage. The exact mechanism of the interaction between NSAIDs and warfarin is unknown, but

may involve enhanced bleeding from NSAID-induced gastrointestinal ulceration or an additive effect of anticoagulation by warfarin and inhibition of platelet function by NSAIDs.

There are isolated reports of suspected interaction of topical formulations of diclofenac with oral anticoagulants.

Since systemic absorption of diclofenac from topical application of the gel is very low, such interactions are very unlikely. Systemic reactions are unlikely to occur when the gel is used as recommended. Nevertheless, the possibility of such an interaction should be borne in mind.

Adverse effects

Adverse reactions (Table 1) are ranked under heading of frequency, the most frequent first, using the following convention: common ($\geq 1/100$, $< 1/10$); uncommon ($\geq 1/1,000$, $< 1/100$); rare ($\geq 1/10,000$, $< 1/1,000$); very rare ($< 1/10,000$), including isolated reports. Within each frequency, adverse effects are presented in order of decreasing seriousness.

Table 1

Infections and infestations	
Very rare:	Rash pustular.
Immune system disorders	
Very rare:	Hypersensitivity (including urticaria), angioedema.
Respiratory, thoracic and mediastinal disorders	
Very rare:	Asthma.
Skin and subcutaneous tissue disorders	
Common:	dermatitis (including contact dermatitis), Rash, erythema, eczema, pruritus
Rare:	Dermatitis bullous.
Very rare:	Photosensitivity reaction.

Dosage and administration

Adults and adolescents aged 12 years and over:

Voltaren Emulgel/ Voltaren Osteo Gel is applied locally to the skin 3 or 4 times daily and rubbed in gently. The amount needed depends on the size of the painful site. For example, 2 to 4 g of gel (a quantity ranging in size from a cherry to a walnut) is sufficient to apply to an area of about 400-800 cm². After application, the hands should be washed, unless they are the site being treated.

Voltaren Emulgel/Voltaren Osteo Gel should never be taken by mouth. Voltaren Emulgel/Voltaren Osteo Gel may be used with non-occlusive bandages but should not be used with occlusive dressings. The duration of treatment depends on the indication and the response obtained. The gel should not be used for more than 14 days for soft-tissue injuries or soft tissue rheumatism, or 21 days for osteoarthritis pain, unless recommended by a doctor.

When used without medical prescription, patients should consult their doctor or pharmacist if the condition does not improve within 7 days, or if it gets worse.

Children:

Voltaren Emulgel/Voltaren Osteo Gel is not recommended for use in children below 12 years of age.

Overdosage

Symptoms and signs

The low systemic absorption of topical diclofenac renders overdosage extremely unlikely. However, undesirable effects, similar to those observed following an overdose of Voltaren tablets, can be expected if the gel is inadvertently ingested (1 tube of 100 g contains the equivalent of 1 g diclofenac sodium).

Treatment

In the event of accidental ingestion, resulting in significant systemic side-effects, general therapeutic measures normally adopted to treat poisoning with non-steroidal anti-inflammatory drugs, should be used. For further information on the treatment of overdose, contact the Poisons Information Centre (Australia 131 126; New Zealand 0800 764 766).

Presentation and storage conditions

Voltaren Emulgel: White to practically white, soft, homogeneous, cream-like gel. Packaged in aluminium laminated tubes (LDPE/aluminium/HDPE) in pack sizes of 20g, 50g, 100g and 120 g
Voltaren Osteo Gel: White to practically white, soft, homogeneous, cream-like gel. Packaged in aluminium laminated tubes (LDPE/aluminium/HDPE) in pack sizes of 75 g, 150 g.

Store below 30°C.

Name and address of the sponsor

GlaxoSmithKline Consumer Healthcare
82 Hughes Ave, Ermington NSW 2115, Australia
FREECALL Australia: 1800 028 533
® = Registered Trade Mark

Poison Schedule of the medicine

Unscheduled

Date of first inclusion on the Australian Register of Therapeutic Goods (the ARTG)

Voltaren Emulgel: 21st April 1994
Voltaren Osteo Gel: 17th September 2010

Date of most recent amendment

Date of most recent amendment: July 2016