

- ▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. You can report side effects to your doctor, or directly at www.tga.gov.au/reporting-problems.

SKYRIZI®

risankizumab

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about SKYRIZI. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you using SKYRIZI against the benefits they expect it will have for you.

If you have any concerns about using this medicine, ask your doctor or pharmacist.

Read this leaflet carefully before you start using this medicine. It includes important information on the safe and effective use of this medicine.

Keep this leaflet with the medicine.

You may need to read it again.

What SKYRIZI is used for

SKYRIZI is used to treat adults with moderate to severe plaque psoriasis, an inflammatory condition affecting the skin and nails.

SKYRIZI contains the active ingredient risankizumab, which is a type of protein called a monoclonal antibody. It works by stopping a protein in the body called IL-23 which causes inflammation. SKYRIZI reduces the inflammation which improves skin clearance and the appearance of the nails. It also reduces symptoms of psoriasis such as burning, itching, pain, redness and scaling.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

Your doctor may have prescribed it for another reason.

This medicine is not expected to affect your ability to drive a car or operate machinery.

Before you use SKYRIZI

When you must not use it

Do not use this medicine if you have an allergy to risankizumab or any of the other ingredients listed at the end of this leaflet.

Do not use this medicine if you have an active infection which your doctor thinks is important.

Do not use this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start using this medicine, talk to your doctor.

Before you start to use it

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor, pharmacist or nurse before and during use of SKYRIZI if you have or have had any of the following medical conditions:

- you currently have an infection or if you have an infection that keeps coming back.
- you have tuberculosis (TB).
- you have recently received or plan to receive an immunisation (vaccine). You should not be given certain types of vaccines while using SKYRIZI.

Tell your doctor if you are pregnant or plan to become pregnant or are breast-feeding.

Your doctor can discuss with you the risks and benefits involved.

If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before using this medicine. This is because it is not known how this medicine will affect the baby.

If you are breast-feeding or are planning to breast-feed, talk to your doctor before using this medicine.

If you have not told your doctor about any of the above, tell him/her before you start using SKYRIZI.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

How to use SKYRIZI

Follow all directions given to you by your doctor, pharmacist or nurse carefully.

If you do not understand the instructions on the box, ask your doctor, pharmacist or nurse for help.

How much to use

SKYRIZI is given as 2 injections under your skin (called 'subcutaneous injections').

The dose is 150 mg given as two 75 mg injections.

- 1st dose = 150 mg (two 75 mg injections) as directed by your doctor, pharmacist or nurse
- 2nd dose = 150mg (two 75 mg injections) 4 weeks after the 1st dose
- Further doses = 150mg (two 75 mg injections) every 12 weeks starting after the 2nd dose


How to use it

You and your doctor, pharmacist or nurse will decide if you should inject SKYRIZI yourself. Do not inject yourself with this medicine unless you have been trained by your doctor, pharmacist or nurse. A caregiver may also give your injections after training.

Special disposal container can be obtained from your local pharmacy.

Read the 'Instructions for Use' before injecting SKYRIZI yourself.

Instructions for Use



Important information to know before you inject SKYRIZI.

- SKYRIZI is for single use in one patient only. Discard any residue.
- Receive training on how to inject SKYRIZI before giving an injection. Talk to your doctor, pharmacist or nurse if you need help.
- Mark the dates on your calendar so you know when to next use SKYRIZI.
- Keep SKYRIZI in the original carton to protect from light until it is time to use it.
- **Do not inject if the liquid is cloudy or contains flakes or large particles. The liquid should look clear to slightly yellow and may contain tiny white or clear particles.**
- **Do not use if the expiration date has passed.**
- **Do not use if the liquid has been frozen (even if thawed).**
- **Do not use if the syringe has been dropped or damaged.**
- **Do not use if the syringe tray cover is broken or missing. Return this medicine to the pharmacy.**

- **Do not remove the needle cover until just before the injection.**

For a more comfortable injection:

Take the carton out of the refrigerator and leave it at room temperature, out of direct sunlight, for 15 to 30 minutes before injecting.

- **Do not remove the syringes from the carton until ready to inject.**
- **Do not warm SKYRIZI in any other way. For example, do not warm it in a microwave or in hot water.**

Follow the steps below each time you use SKYRIZI.

STEP 1



Place the following on a clean, flat surface:

- 2 pre-filled syringes and 2 alcohol pads (provided).
- 2 cotton balls or gauze pads (not provided).

- Special disposal container (not provided).


Wash and dry your hands.

Start with one syringe for the first injection.

For a full dose, 2 injections are required, one after the other.

STEP 2

Areas to inject



Areas to inject

Choose from these 3 areas to inject:

- Front of left thigh.
- Front of right thigh.
- Belly (abdomen) at least 5 cm from the belly button (navel).


For the second syringe, inject at least 3 cm away from the first injection.

- **Do not inject into the same place.**

Before each injection, wipe where you will inject in a circular motion with an alcohol pad.

- **Do not inject through clothes.**
- **Do not inject into skin that is sore, bruised, red, hard, scarred or has stretch marks.**
- **Do not inject into areas affected by psoriasis.**

STEP 3




Hold the syringe with covered needle pointing down, as shown.

Check the liquid in the syringe.

- It is normal to see bubbles in the window.

- The liquid should look clear to slightly yellow and may contain tiny white or clear particles.
- **Do not use if the liquid is cloudy or contains flakes or large particles.**


STEP 4



Removing the needle cover:

- Hold the syringe in one hand.
- With the other hand, gently pull the needle cover straight off.
- You may see a drop of liquid at the end of the needle. This is normal.
- Throw away the needle cover.
- **Do not touch the needle with your fingers or let the needle touch anything.**

STEP 5




Hold the body of the syringe in one hand between the thumb and index finger, like you would a pencil.

Gently pinch the area of cleaned skin with your other hand and hold it firmly.

Insert the needle all the way into the skin at about a 45-degree angle using a quick, short movement. Keep the syringe steady at the same angle.

STEP 6



Needle guard

Slowly push the plunger all the way in until all of the liquid is injected.

Pull the needle out of the skin while keeping the syringe at the same angle.

Stop pinching the skin around your injection site.


Slowly take your thumb off the plunger. The needle will then be covered by the needle guard.

- The needle guard will not activate unless all the liquid is injected.
- Speak to your doctor, pharmacist or nurse if you think you may have not given a full dose.

Press a cotton ball or gauze pad where you have injected and hold for 10 seconds.

Do not rub the skin where you have injected. You may have slight bleeding from where you injected. This is normal.

STEP 7




2 Injections Required

For a full dose, two injections are needed, one after the other.

- Repeat Steps 2 through 6 with the second syringe.
- Inject the second syringe straight after the first injection.

STEP 8



Throw away used syringes in a special disposal container.

- **Do not throw away used syringes in the household waste.**
- Your doctor, pharmacist or nurse will tell you how to return the full special disposal container.

How long to use it

Do not stop using SKYRIZI without talking to your doctor first. If you stop treatment, your symptoms may come back.

If you forget to use it

If you forget to use SKYRIZI, inject a dose as soon as you remember. Talk to your doctor or pharmacist if you are not sure what to do.

If you use too much (overdose)

If you have used more SKYRIZI than you should, or the dose has been given sooner than prescribed, talk to your doctor.

Immediately telephone your doctor or the Poisons Information Centre (telephone 13 11 26) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have used too much SKYRIZI. Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

While you are using SKYRIZI

Things you must do

If you are about to be started on any new medicine, remind your doctor and pharmacist that you are using SKYRIZI.

Tell any other doctors, dentists, and pharmacists who treat you that you are using this medicine.

If you become pregnant while using this medicine, tell your doctor immediately.

Things you must not do

You should not receive a live vaccine while taking SKYRIZI. Talk to your doctor, pharmacist or nurse before receiving any vaccination while taking SKYRIZI. Do not use SKYRIZI to treat any other complaints unless your doctor tells you to.

Do not give your medicine to anyone else, even if they have the same condition as you.

Do not stop using your medicine or lower the dosage without checking with your doctor.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are using SKYRIZI.

This medicine helps most people, but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Do not be alarmed by the following lists of side effects. You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Talk to your doctor or get medical help as soon as possible if you have symptoms of serious infection such as:

- Fever, flu-like symptoms, night sweats.
- Feeling tired or short of breath, cough which will not go away.
- Warm, red and painful skin, or a painful skin rash with blisters.

Your doctor will decide if you can keep using SKYRIZI.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- Upper respiratory infections with symptoms such as a sore throat and stuffy nose
- Feeling tired
- Fungal skin infection
- Injection site reactions
- Headache
- Small raised red bumps on the skin.

The above list includes the more common side effects of SKYRIZI.

Tell your doctor or pharmacist if you notice anything that is making you feel unwell.

Other side effects not listed above may also occur in some people.

After using SKYRIZI

Storage

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiration date which is stated on the syringe label and outer carton after 'EXP'.

Store SKYRIZI at 2°C to 8°C. Refrigerate. Do not freeze.

Keep the pre-filled syringes in the original carton in order to protect from light.

The liquid should look clear to slightly yellow. The liquid may contain tiny white or clear particles. Do not use if the liquid is cloudy or contains flakes or large particles.

Disposal

After injecting SKYRIZI, immediately throw away the used syringes in a special container as instructed by your doctor, nurse or pharmacist.

If your doctor tells you to stop using this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

Product description

What it looks like

SKYRIZI is a clear and colourless to slightly yellow liquid in a pre-filled syringe with needle guard.

Ingredients

SKYRIZI contains 75 mg of risankizumab as the active ingredient in 0.83 mL solution in each pre-filled syringe.

The other ingredients include:

- Sodium succinate hexahydrate
- Succinic acid
- Sorbitol
- Polysorbate 20
- Water for injections

Distributor

SKYRIZI is distributed in Australia by:

AbbVie Pty Ltd

241 O'Riordan Street

Mascot NSW 2020

Australia

SKYRIZI is distributed in New Zealand by:

AbbVie Limited

6th Floor, 156 158 Victoria Street

Wellington 6011

New Zealand

This leaflet was prepared in

July 2019

AUST R 304226

Version 01