


ELOCON CREAM, OINTMENT AND LOTION PRODUCT INFORMATION

NAME OF THE MEDICINE

Mometasone furoate 0.1% (1 mg/g)

Chemical structure:



Mometasone furoate is 9 α ,21-dichloro-11 β ,17-dihydroxy-16 α -methylpregna-1,4-diene-3,20-dione 17-(2-furoate). The empirical formula is C₂₇H₃₀Cl₂O₆. MW: 521.4.

DESCRIPTION

Mometasone furoate is a white to off-white powder practically insoluble in water, slightly soluble in octanol and moderately soluble in ethyl alcohol.

Each gram of ELOCON Cream contains mometasone furoate 1mg in a cream base of soft white paraffin, hexylene glycol, aluminium starch octenylsuccinate, propylene glycol monostearate, stearyl alcohol, cetareth-20, white beeswax, purified water, titanium dioxide and phosphoric acid.

Each gram of ELOCON Ointment contains mometasone furoate 1mg in an ointment base of soft white paraffin, hexylene glycol, white beeswax, purified water, propylene glycol monostearate and phosphoric acid.

Each gram of ELOCON Lotion contains mometasone furoate 1mg in a lotion base of isopropyl alcohol, propylene glycol, hydroxypropylcellulose, sodium phosphate monobasic dihydrate, phosphoric acid and purified water.

PHARMACOLOGY

Mometasone furoate is a synthetic corticosteroid, exhibiting anti-inflammatory, antipruritic and vasoconstrictive properties.

In laboratory animals, mometasone furoate exhibits potent topical anti-inflammatory activity but approximately half of the suppressive effect on the HPA (hypothalamic-pituitary-adrenal) axis when compared with equivalent doses of betamethasone valerate. The topical to systemic potency ratio of mometasone furoate is approximately 3 to 10 times that of betamethasone valerate in animal studies.

Pharmacokinetics

Following topical application of radio-labelled mometasone furoate in animals, systemic absorption was minimal in all species studied, ranging from approximately 2% in dogs to 11% in rabbits over a 5 to 7 day period.

The percutaneous absorption of ELOCON was evaluated in healthy volunteers receiving a single application of radio-labelled mometasone furoate cream 0.1% which remained on intact skin for eight hours. Based on the radioactivity excreted in the urine and faeces during the five day study period, approximately 0.4% of the applied dose was absorbed systemically. In a similar study conducted using the ointment formulation, approximately 0.7% of the applied dose was absorbed systemically.

Inflammation and/or other disease processes in the skin may increase percutaneous absorption. Occlusive dressings substantially increase the percutaneous absorption of topical corticosteroids. As Elocon is applied topically and only low concentrations of radioactivity are detected in plasma, specific bioavailability studies have not been conducted for mometasone furoate. Since plasma levels of radiolabelled product are very low, metabolism in humans has not been studied.

INDICATIONS

ELOCON Cream, Ointment and Lotion are indicated for short-term (up to four (4) continuous weeks) relief of inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses, such as psoriasis and atopic dermatitis.

ELOCON Lotion is also suitable for short-term use for scalp psoriasis and seborrhoeic dermatitis.

CONTRAINDICATIONS

ELOCON Cream, Ointment and Lotion are contraindicated in patients who are hypersensitive to mometasone furoate or to other corticosteroids. Like other topical corticosteroids, ELOCON is contraindicated in most viral infections of the skin, tuberculosis, acne rosacea, perioral dermatitis, fungal skin infections and ulcerative conditions.

PRECAUTIONS

If irritation or sensitisation develops with the use of ELOCON Cream, Ointment or Lotion treatment should be discontinued and appropriate therapy instituted.

In the presence of an infection, use of an appropriate antifungal or antibacterial agent should be instituted. If a favourable response does not occur promptly, the corticosteroid should be discontinued until the infection is controlled adequately.

Any of the side effects that have been reported following systemic use of corticosteroids, including adrenal suppression, may also occur with topical corticosteroids, especially in infants and children.

Systemic absorption of topical corticosteroids will be increased if extensive body surface areas are treated, if the occlusive technique is used, if used in areas where the epidermal barrier is disrupted or if used long-term. Suitable precautions should be taken to ensure application sites are not occluded, particularly in infants and children.

In infants, plastic pants and napkins may act as occlusive dressings and increase absorption. Paediatric patients may demonstrate greater susceptibility to topical corticosteroid-induced HPA axis suppression and Cushing's syndrome than adults because of a larger skin surface area to body weight ratio. Use of topical corticosteroids in children should be limited to the least amount required for a therapeutic effect. Chronic corticosteroid therapy may interfere with growth and development of children.

ELOCON Cream, Ointment and Lotion are not for ophthalmic use.

Use in Pregnancy (Category B3)

Category B3: Drugs which have been taken by only a limited number of pregnant women and women of childbearing age, without an increase in the frequency of malformation or other direct or indirect harmful effects on the human foetus having been observed. Studies in animals have shown evidence of an increased occurrence of foetal damage, the significance of which is considered uncertain in humans.

Corticosteroids are generally teratogenic in laboratory animals when administered systemically at relatively low dosage levels. Similarly mometasone furoate has been shown to be teratogenic after dermal application to animals. At doses greater than 0.3 mg/kg in rats and at all dose levels tested in rabbits (0.15 mg/kg and 0.3 mg/kg), sequelae typical of other topical corticosteroids resulted. There are no adequate and well controlled studies of the teratogenic effects of corticosteroids in pregnant women. Therefore, topical corticosteroids should be used during pregnancy only if the potential benefit justifies the potential risk to the foetus. Drugs of this class should not be used on pregnant patients in large amounts or for prolonged periods of time.

Use in Lactation

It is not known whether topical administration of corticosteroids could result in sufficient systemic absorption to produce detectable quantities in breast milk. Systemically administered corticosteroids are secreted into breast milk in quantities not likely to have a deleterious effect on the infant. Nevertheless, a decision should be made whether breast-feeding should be discontinued or ELOCON Cream, Ointment or Lotion be discontinued, taking into account the importance of the drug to the mother.

ADVERSE EFFECTS

ELOCON Cream, Ointment and Lotion are generally well tolerated. Pruritis, burning, tingling/stinging, signs of skin atrophy and acneiform reaction have been reported in less than 5% of patients.

Other local adverse reactions reported in less than 1% of patients include erythema, furunculosis, dermatitis, abscess, aggravated allergy, increased lesion size, disease exacerbation, paraesthesia, dry skin, pimples, folliculitis and papular and pustular formation.

The following local adverse reactions have been reported infrequently with the use of other topical corticosteroids: irritation, hypertrichosis, hypopigmentation, perioral

dermatitis, allergic contact dermatitis, maceration of the skin, secondary infection, striae and miliaria.

DOSAGE AND ADMINISTRATION

A thin film of ELOCON Cream or Ointment should be applied to the affected skin areas once daily. ELOCON Cream is suitable for moist lesions; the ointment should be used for dry, scaling and fissured lesions.

A few drops of ELOCON Lotion should be applied to affected skin areas including scalp sites once daily; massage gently and thoroughly until the medication disappears.

OVERDOSAGE

Excessive, prolonged use of topical corticosteroids can suppress pituitary-adrenal function resulting in secondary adrenal insufficiency.

Treatment: Appropriate symptomatic treatment is indicated. Acute hypercorticoid symptoms are virtually reversible. Treat electrolyte imbalance, if necessary. In cases of chronic toxicity, slow withdrawal of corticosteroids is advised.

PRESENTATION AND STORAGE CONDITIONS

ELOCON Ointment: 15 g, 45 g and 50 g tubes

ELOCON Cream: 15 g, 45 g and 50 g tubes

ELOCON Lotion: 10mL*, 15mL*, 20mL*, 30mL, 50mL* and 100mL* bottles

* not currently available in Australia

Cream, Ointment and Lotion: Store below 25°C.

NAME AND ADDRESS OF THE SPONSOR

Merck Sharp & Dohme (Australia) Pty Limited
Level 1, Building A, 26 Talavera Road
Macquarie Park, NSW 2113
Australia

Merck Sharp & Dohme (New Zealand) Ltd
P O Box 99 851
Newmarket
Auckland 1149
New Zealand

POISONS SCHEDULE OF THE MEDICINE

Schedule 4 – Prescription Only Medicines

DATE OF APPROVAL

This product information was approved by the Therapeutic Goods Administration on 23rd March 2011.

Date of most recent amendment: 7 July 2014