

PRODUCT INFORMATION


REMERON SolTab®

NAME OF THE MEDICINE

Non-proprietary Name

Mirtazapine

Chemical Structure



Mirtazapine - INN name

Molecular Formula C₁₇H₁₉N₃

Molecular Mass 265.36

Chemical name: (±)-1,2,3,4,10,14b-hexahydro-2-methyl-pyrazino[2,1-a]pyrido[2,3-c][2]benzazepine.

CAS Number

CAS No. 61337-67-5

DESCRIPTION

Mirtazapine is a tetracyclic piperazinoazepine analogue of mianserin, a chemical structure unrelated to tricyclic antidepressants, monoamine oxidase inhibitors or selective serotonin reuptake inhibitors. Mirtazapine is a white to creamy white crystalline powder which is slightly soluble in water.

Qualitative and Quantitative Composition

Each tablet contains 15 mg, 30 mg or 45 mg of mirtazapine.

List of excipients

Remeron SolTab 15 mg, 30 mg and 45 mg tablets contain: sucrose, starch, hypromellose, povidone, Eudragit E100, magnesium stearate, mannitol, crospovidone, sodium bicarbonate, citric acid anhydrous, cellulose microcrystalline, aspartame, and orange flavour SN 027512.

PHARMACOLOGY

Pharmacodynamics

Remeron SolTab is an antidepressant, which can be given as treatment for episodes of major depression. Mirtazapine is an antagonist of central α_2 -auto and hetero-adrenoceptors which causes an increase in both noradrenaline and serotonin release. The effect of released serotonin is exerted

specifically via 5-HT₁ type receptors, because 5-HT₂ and 5-HT₃ type receptors are specifically blocked by mirtazapine. Mirtazapine is accordingly a noradrenergic and specific serotonergic antidepressant. The α_2 , 5-HT₂ and 5-HT₃ antagonistic effects all contribute to the antidepressant profile of mirtazapine. The presentation of mirtazapine is as a racemate. The two enantiomers contribute differently to its pharmacological profile. The α_2 and 5-HT₂ receptor blocking activity is contained in the (S)+ enantiomer, whereas the 5-HT₃ receptor blocking activity is contained in the (R)- enantiomer. The presence of both enantiomers is therefore considered to be essential for the antidepressant activity of mirtazapine. In one study, there was no efficacy difference indicated between the two enantiomers, despite their different receptor affinities.

Mirtazapine is generally well tolerated. The histamine H₁-antagonistic activity of mirtazapine may cause a degree of sedation in the first weeks of treatment. It has practically no anticholinergic activity. Mirtazapine has been associated with acute postural hypotension in healthy volunteer studies but this occurred rarely in patient studies (see **ADVERSE EFFECTS**).

Pharmacokinetics

Absorption

After oral administration of Remeron SolTab tablets, the active substance mirtazapine is rapidly and well absorbed (bioavailability \cong 50%), reaching peak plasma levels after about 2 hours. Food intake has no clinically significant influence on the pharmacokinetics of mirtazapine.

Distribution

Binding of mirtazapine to plasma proteins is approx. 85%. The half-life of elimination ranged from 20-40 hours; longer half-lives, up to 65 hours, have occasionally been recorded and shorter half-lives have been seen in young men. The half-life of elimination is sufficient to justify once-a-day dosing. Steady state is reached after 3-6 days, after which there is no further accumulation. Mirtazapine displays linear pharmacokinetics within the recommended dose range.

Metabolism

In vitro data from human liver microsomes indicate that cytochrome P450 enzymes CYP 2D6 and CYP 1A2 are involved in the formation of the 8-hydroxy metabolite of mirtazapine, whereas CYP 3A4 is considered to be responsible for the formation of the N-demethyl and N-oxide metabolites.

The presentation of mirtazapine is as a racemate. It is not known whether first pass extraction of the drug is stereoselective but it is known that the clearance of the two enantiomers is by different metabolic processes.

Elimination

Mirtazapine is extensively metabolized and its metabolites are eliminated via the urine and faeces within four days. Major pathways of biotransformation are demethylation and oxidation, followed by conjugation. The demethyl metabolite is pharmacologically active and appears to have the same pharmacokinetic profile as the parent compound.

Special populations

Renal and/or hepatic impairment:

The clearance of mirtazapine may be decreased as a result of renal or hepatic insufficiency. Mirtazapine is substantially excreted by the kidney (75%) and the risk of decreased clearance of this drug is greater in patients with impaired renal function, refer to **DOSAGE AND ADMINISTRATION**.

Geriatric:

The recommended dosage regimen is the same as for adults. Increases should be monitored carefully, refer to **DOSAGE AND ADMINISTRATION**.

Children and adolescents

The safety and effectiveness of mirtazapine has not been established in children and adolescents and therefore should not be prescribed in these patient groups, refer **PRECAUTIONS**.

Sex:

The half-life of elimination of mirtazapine ranged from 20-40 hours, longer half-lives up to 65 hours have occasionally been recorded and shorter half-lives have been seen in young men.

Race:

There is no information available regarding the effect of race on the pharmacokinetics of mirtazapine.

CLINICAL TRIALS

Several placebo-controlled double-blind studies have demonstrated that mirtazapine is statistically significantly more effective than placebo in the short term treatment of a major depressive episode; the efficacy is maintained during continuation treatment with mirtazapine.

Active controlled studies

The efficacy of mirtazapine has been found to be comparable to several standard antidepressant agents (amitriptyline, doxepin, clomipramine). In addition, eleven 6 or 8 week studies and a 24 week study have been performed in moderately to severely depressed patients in which efficacy and tolerability of mirtazapine were compared to SSRIs (4 vs fluoxetine, 3 vs paroxetine, 2 vs sertraline, 2 vs fluvoxamine and 1 vs citalopram). The primary efficacy parameters in these studies were:

- change from baseline on HAM-D total score (Hamilton depression rating scale, 17 items). 7 studies.
- proportion or number of HAM-D 50% responders. 3 studies.
- change from baseline on MADRAS total score (Montgomery-Asberg depression rating scale, 10 items). 1 study.
- VAMRS 6 items (Visual Analogue Mood Rating Scale). 1 study. Change in HAM-D (12 items) total score was a secondary parameter in this study.

On an intention-to-treat basis, a total of 1402 patients were treated with mirtazapine and 1405 patients were treated with the comparator. In all 12 studies, mirtazapine proved to be at least comparable in efficacy to the SSRIs. In 11 of these studies, statistically significant greater reductions in HAM-D or MADRS total scores and more responders were observed in the mirtazapine groups at one or more timepoints in the first 4 weeks.

A meta-analysis of these 12 studies provides further comparison of the onset of efficacy of mirtazapine relative to the SSRIs studied. The primary efficacy parameter for this meta-analysis was time to first 50% reduction on recalculated HAM-D total score (17 items) or recalculated MADRS total score (10 items). There were also a number of secondary parameters which are identified in Tables 1 and 2. Table 1 provides an analysis of the relative event rates (estimated hazard ratios) for various depression parameters limited to the first 3 treatment weeks for the occurrence of the event and the entire 6-8 week study period to define whether the event was sustained or not. The increased hazard ratios demonstrate that the probability at any time t of first response (50% or more score reduction), remission, sustained response or sustained remission was consistently and significantly greater among mirtazapine-treated than SSRI-treated patients, indicating an earlier onset of efficacy. The statistically earlier onset of action observed with mirtazapine may not necessarily translate in to a meaningful clinical benefit for an individual patient. Table 2 presents the proportions of HAM-D responders and HAM-D/MADRS remitters at the various time points during treatment. At most time points there were significantly more responders and remitters among mirtazapine-treated patients than among SSRI-treated patients.

Table 1.

Parameter * primary ** secondary	Estimated hazard ratio mirtazapine relative to SSRI.	95% confidence interval	p-value
first day 50% or more reduction in HAM-D/MADRAS total score *	1.49	1.32 - 1.68	≤ 0.001
first day 50% reduction on HAM-D Bech depression factor **	1.20	1.06 – 1.37	0.005
day of 50% or more sustained reduction in HAM-D/MADRAS total score **	1.58	1.37 – 1.82	≤ 0.001
day of 50% sustained reduction on HAM-D Bech depression factor **	1.22	1.05 – 1.42	0.010
first day remission on HAM-D total score (≤ 7) or MADRAS total score (≤ 12) **	1.67	1.40 – 2.00	≤ 0.001
day of sustained remission on HAM-D total score (≤ 7) or MADRAS total score (≤ 12) **	1.68	1.35 – 2.09	≤ 0.001

Table 2.

Parameter * primary ** secondary	Cumulative probability mirtazapine (%) v SSRI (%) (Estimated Difference (%) between mirtazapine and SSRI adjusted for trial) (p value)				
	Week 1	Week 2	Week 4	Week 6	EP***
HAM-D responders * (subjects where score is reduced by ≥ 50%)	11.5 v 7.0 (4.4) (≤0.001)	29.5 v 20.8 (8.6) (≤0.001)	50.4 v 40.7 (9.7) (≤0.001)	60.2 v 55.1 (5.2) (0.013)	61.5 v 57.1 (4.4) 0.023
MADRS responders * (subjects where score is reduced by ≥ 50%)	6.6 v 4.7 (1.9) (0.205)	28.8 v 20.1 (8.6) (0.002)	51.6 v 49.0 (2.5) (0.455)	65.3 v 65.3 (-0.1) (0.967)	72.4 v 73.2 (-0.9) (0.768)
HAM-D or MADRS responders * (subjects where score is reduced by ≥ 50%)	10.5 v 6.6 (3.9) (≤0.001)	28.7 v 20.3 (8.3) (≤0.001)	51.5 v 42.2 (9.3) (≤0.001)	61.9 v 57.4 (4.6) (0.018)	63.8 v 60.0 (3.7) (0.038)
HAM-D or MADRAS remitters ** (HAM-D ≤7 MADRS ≤12)	3.4 v 1.8 (1.6) (0.008)	11.8 v 6.9 (4.9) (≤0.001)	28.6 v 21.8 (6.8) (≤0.001)	38.8 v 34.7 (4.1) (0.028)	42.7 v 39.9 (2.7) (0.126)

The shaded cells indicate statistical significance in the result.

*** EP – Endpoint analysis consists of results from week 6 assessments of the 6 week studies and week 8 assessments of the 8 week studies.

Some secondary parameter results have been excluded from Table 2. These were number of:

- 50% Bech responders
- 50% HAM-D Factor I 'anxiety/somatisation' responders
- 50% HAM-D Factor V 'retardation' responders
- 50% HAM-D Factor VI 'sleep disturbance' responders
- HAM-D item 'depressed mood' responders (=0 or <2)
- HAM-D item 'suicide or MADRS item 'suicidal thoughts' (=0 or <2)

Statistically significant differences favouring mirtazapine were observed for HAM-D factors V and VI at week 1 to 6 timepoints. Statistically significant differences favouring mirtazapine were observed for HAM-D factor I at week 1 to 4 timepoints. A statistically significant difference was observed in favour of mirtazapine for Bech responders at the week 2 timepoint. There were no other statistically significant differences.

An eight-week comparative study was performed to compare the antidepressant efficacy and tolerability of mirtazapine and venlafaxine in the treatment of 157 hospitalised patients with severe depression with melancholic features (HAM-D total score > 25). In this study, mirtazapine and venlafaxine were equally effective in reducing symptoms of depression and improving quality of life during treatment.

Elderly

The efficacy and tolerability in elderly patients was investigated in three randomised controlled trials. In two six-week trials with a total of 270 patients aged over 55 years (mean age 70 and 62 years respectively), mirtazapine was at least as effective as amitriptyline and all treatments were well tolerated. In an eight-week study in 255 patients aged 65 and over (mean age 72 years) comparing mirtazapine with paroxetine, mean HAM-D scores were similar at end-point but lower for mirtazapine in the first 3 weeks, although only at day 14 was the difference statistically significant. Total discontinuation rates were similar (22.7% for mirtazapine versus 31.0% for paroxetine), although discontinuation due to adverse events was lower with mirtazapine than paroxetine (14.8% versus 26.2%) and discontinuation due to lack of efficacy higher (3.9% versus 0%).

INDICATIONS

Treatment of major depression

CONTRAINDICATIONS

- Hypersensitivity to mirtazapine or to any of the excipients.
- MAO Inhibitors as concomitant therapy. It is recommended that Remeron SolTab not be used in combination with MAO Inhibitors, or within 14 days of initiating or discontinuing therapy with a MAO Inhibitor (see **INTERACTIONS WITH OTHER MEDICINES**).

PRECAUTIONS

Clinical worsening and suicide risk

The risk of suicidality (suicidal ideation and suicidal behaviours) is inherent in depression and may persist until significant remission occurs. This risk must be considered in all depressed patients. Patients with depression may experience worsening of their depressive symptoms and/or the emergence of suicidal ideation and/or behaviours whether or not they are taking antidepressant medication, and this risk may persist until significant remission occurs. Suicide is a known risk in depression and certain other psychiatric disorders, and these disorders themselves are the strongest predictors of suicide. As improvement may not occur during the first few weeks or more of treatment, patients should be closely monitored for clinical worsening and suicidality, especially at the beginning of a course of treatment, or at the time of dose changes, either increases or decreases. Consideration should be given to changing the therapeutic regimen, including possibly discontinuing the medication, in patients whose depression is persistently worse or whose emergent suicidality is severe, abrupt in onset, or was not part of the patient's presenting symptoms. Patients (and caregivers of patients) should be alerted about the need to monitor for any worsening of their condition and/or the

emergence of suicidal ideation or behaviour or thoughts of harming themselves and to seek medical advice immediately if these symptoms present. Patients with co-morbid depression associated with other psychiatric disorders being treated with antidepressants should be similarly observed for clinical worsening and suicidality.

Pooled analysis of short-term placebo-controlled trials of antidepressant drugs (SSRIs and others) showed that these drugs increased the risk of suicidal ideation and/or behaviours in children, adolescents, and young adults (aged 18-24 years) with major depressive disorder (MDD) and other psychiatric disorders during the initial treatment (generally the first one to two months). Short-term studies did not show an increase in the risk of suicidality with antidepressants compared to placebo in adults beyond the age of 24 years; there was a reduction with antidepressants compared to placebo in adults aged 65 years and older.

The pooled analyses of placebo-controlled trials in children and adolescents with MDD, obsessive compulsive disorder (OCD), or other psychiatric disorders included a total of 24 short-term trials (4 to 16 weeks) of 9 antidepressant drugs in over 4,400 patients. The pooled analyses of placebo-controlled trials in adults with MDD or other psychiatric disorders included a total of 295 short-term trials (median duration of 2 months) of 11 antidepressant drugs in over 77,000 patients. There was considerable variation in the risk of suicidality among drugs, but a tendency towards an increase in the younger patients for almost all drugs studied. There were differences in absolute risk of suicidality across different indications, with the highest incidence in MDD trials. The risk differences (drug vs placebo), however, were relatively stable within age strata and across indications.

No suicides occurred in any of the paediatric trials. There were few suicides in the adult trials, but the number was not sufficient to reach any conclusion about the effect of antidepressants on suicide. It is unknown whether suicidality risk extends to longer-term use, i.e., beyond several months. However, there is substantial evidence from placebo-controlled maintenance trials in adults with depression that the use of antidepressants can delay the recurrence of depression.

Symptoms of anxiety, agitation, panic attacks, insomnia, irritability, hostility, (aggressiveness), impulsivity, akathisia (psychomotor restlessness), hypomania, and mania, have been reported in adults, adolescents and children being treated with antidepressants for major depressive disorders as well as for other indications, both psychiatric and non-psychiatric. Although a causal link between the emergence of such symptoms and either worsening of depression and/or emergence of suicidality impulses has not been established, there is concern that such symptoms may be precursors of emerging suicidality.

Families and caregivers of children and adolescents being treated with antidepressants for MDD or for any other condition (psychiatric or non-psychiatric) should be informed about the need to monitor these patients for the emergence of agitation, irritability, unusual changes in behaviour, and other symptoms described above, as well as the emergence of suicidality and to report such symptoms immediately to health care providers. It is particularly important that monitoring be undertaken during the initial few months of antidepressant treatment or at times of dose increase or decrease.

Prescriptions for Remeron SolTab should be written for the smallest quantity of tablets consistent with good patient management, in order to reduce the risk of overdose.

Conditions which need supervision

Careful dosing as well as regular and close monitoring is necessary in patients with:

- **Hepatic impairment: Epilepsy and organic brain syndrome, refer to ADVERSE EFFECTS:** Remeron SolTab should be introduced cautiously in patients who have a history of seizures. Treatment should be discontinued in any patient who develops seizures, or where there is an increase in seizure frequency.
- **Renal insufficiency:** Mirtazapine is substantially excreted by the kidney (75%) and the risk of decreased clearance of this drug is greater in patients with impaired renal function. **Cardiac diseases:** Such as conduction disturbances, angina pectoris and recent myocardial infarct, where normal precautions should be taken and concomitant medicines carefully administered.

- **Low blood pressure** and conditions that would predispose patients to hypotension (dehydration, hypovolemia and treatment with antihypertensive medication).
- **Diabetes mellitus:** In patients with diabetes, antidepressants may alter glycemic control. Insulin and/or oral hypoglycemic dosage may need to be adjusted and close monitoring is recommended.

Like with other antidepressants, the following should also be taken into account:

- Worsening of psychotic symptoms can occur when antidepressants are administered to patients with schizophrenia or other psychotic disturbances; paranoid thoughts can be intensified.
- A major depressive episode may be the initial presentation of bipolar disorder. It is generally believed that treating such an episode with an antidepressant alone can increase the likelihood of precipitation of a mixed/manic episode in patients at risk of bipolar disorder. Prior to initiating treatment with an antidepressant, patients should be adequately screened to determine if they are at risk for bipolar disorder; such screening should include a detailed psychiatric history, including a family history of suicide, bipolar disorder and depression. When the depressive phase of the bipolar disorder is being treated, it can transform into the manic phase. Patients with a history of mania/hypomania should be closely monitored. Remeron SolTab should be discontinued in any patient entering a manic phase.
- Care should be taken in patients with micturition disturbances like prostate hypertrophy (although problems are not to be expected because mirtazapine possesses only very weak anticholinergic activity).
- Acute narrow-angle glaucoma and increased intra-ocular pressure (however mirtazapine has weak anticholinergic activity).
- Akathisia/psychomotor restlessness: The use of antidepressants have been associated with the development of akathisia, characterized by a subjectively unpleasant or distressing restlessness and need to move often accompanied by an inability to sit or stand still. This is most likely to occur within the first few weeks of treatment. In patients who develop these symptoms, increasing the dose may be detrimental.
- The effect of mirtazapine on QTc interval was assessed in a randomized, placebo and moxifloxacin controlled clinical trial involving 54 healthy volunteers using exposure response analysis. This trial revealed that both 45 mg (therapeutic) and 75 mg (supratherapeutic) doses of mirtazapine did not affect the QTc interval to a clinically meaningful extent. During the postmarketing use of mirtazapine, cases of QT prolongation, Torsades de Pointes, ventricular tachycardia, and sudden death, have been reported. The majority of reports occurred in association with overdose or in patients with other risk factors for QT prolongation, including concomitant use of QTc prolonging medicines (see **INTERACTION WITH OTHER MEDICINES** and **OVERDOSAGE** sections). Caution should be exercised when Remeron SolTab is prescribed in patients with known cardiovascular disease or family history of QT prolongation, and in concomitant use with other medicinal products thought to prolong the QTc interval.
- Remeron SolTab is not addictive. Post-marketing experience shows that abrupt termination of treatment after long term administration may sometimes result in withdrawal symptoms. The majority of withdrawal reactions are mild and self limiting. Among the various reported withdrawal symptoms, dizziness, agitation, anxiety, headache and nausea are the most frequently reported. Even though they have been reported as withdrawal symptoms, it should be realised that these symptoms may be related to underlying disease. As advised in **DOSAGE AND ADMINISTRATION**, it is recommended to discontinue treatment with Remeron SolTab gradually.

Jaundice

Treatment should be discontinued if jaundice occurs.

Hyponatremia

Hyponatremia has been reported very rarely with the use of mirtazapine. Caution should be exercised in patients at risk, such as elderly patients or patients concomitantly treated with medications known to cause hyponatremia.

Serotonin syndrome

Development of serotonin syndrome may occur in association with treatment with SSRIs and SNRIs, particularly when given in combination with MAO-Is (see **CONTRAINDICATIONS and INTERACTION WITH OTHER MEDICINES**) or other serotonergic agents (see **INTERACTION WITH OTHER MEDICINES**). Symptoms and signs of serotonin syndrome include rapid onset of neuromuscular

excitation (hyperreflexia, incoordination, myoclonus, tremor), altered mental status (confusion, agitation, hypomania) and autonomic dysfunction (diaphoresis, diarrhoea, fever, shivering and rapidly fluctuating vital signs). Treatment with Remeron SolTab should be discontinued if such events occur and supportive symptomatic treatment initiated. From post marketing experience it appears that serotonin syndrome occurs very rarely in patients treated with mirtazapine alone (see **ADVERSE EFFECTS**).

Phenylketonurics

Phenylketonuric patients should be informed the Remeron SolTab contains aspartame, a source of phenylalanine. Each tablet with 15 mg, 30 mg and 45 mg mirtazapine corresponds to 2.6 mg, 5.2 mg and 7.8 mg phenylalanine respectively. It may be harmful for patients with phenylketonuria.

Remeron SolTab contains sugar spheres containing sucrose. Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrose-isomaltase insufficiency should not take this medicine.

Elderly patients

Elderly patients are often more sensitive, especially with regard to the undesirable effects of antidepressants. During clinical research with mirtazapine, undesirable effects have not been reported more often in elderly patients than in other age groups (refer to **CLINICAL TRIALS** and **DOSAGE AND ADMINISTRATION**).

Use in children and adolescents (<18 years of age)

Remeron SolTab should not be used to treat children and adolescents under the age of 18 years. Suicide-related behaviors (suicide attempt and suicidal thoughts), and hostility (predominantly aggression, oppositional behavior and anger) were more frequently observed in clinical trials among children and adolescents treated with antidepressants compared to those treated with placebo. If, based on clinical need, a decision to treat is nevertheless taken; the patient should be carefully monitored for the appearance of suicidal symptoms. In addition, long-term safety data in children and adolescents concerning growth, maturation and cognitive and behavioral development are lacking.

Neutropenia, Agranulocytosis

Bone marrow depression, usually presenting as granulocytopenia or agranulocytosis, has been reported during treatment with mirtazapine. The symptoms mostly appear after 2-6 weeks of treatment. The bone marrow depression is, in general, reversible after termination of treatment. However in very rare cases agranulocytosis can be fatal. Reversible agranulocytosis has been reported as a rare occurrence in clinical studies with mirtazapine. In the post-marketing period with Remeron SolTab very rare cases of agranulocytosis have been reported, mostly reversible, but in some cases fatal. All fatal cases concerned patients over 65 years. Post-marketing data indicate that the rate of occurrence of agranulocytosis and agranulocytosis-like disorders (whether or not causally related) amongst mirtazapine users is no greater than that in the background population. One should be alert for symptoms like fever, sore throat, stomatitis or other signs of infections. If such symptoms occur the treatment should be stopped and blood counts taken.

Carcinogenicity

An eighteen month carcinogenicity study in mice showed an increase in the development of hepatic tumours in males after mirtazapine treatment at oral doses of 20 mg/kg/day and above. In a two year carcinogenicity study in rats, oral doses of mirtazapine greater than 20 mg/kg/day were associated in males with an increased incidence of thyroid follicular cell adenomas and carcinomas.

Mutagenicity

Since the only tumours found in carcinogenicity studies with mice and rats were considered to be species-specific, non-genotoxic responses associated with long term treatment with hepatic enzyme inducers, mirtazapine is not expected to possess carcinogenic potential at therapeutic dosages in the clinic. Mirtazapine was not genotoxic in a series of tests for gene mutation and chromosomal and DNA damage.

Impairment of fertility

In a fertility study in rats, mirtazapine was given at doses up to 100 mg/kg [ca. 20 times the recommended human dose of 45 mg on a mg/m² basis]. The drug did not affect mating and

conception, but oestrus cycling was disrupted at doses that were 3 or more times the recommended human dose of 45 mg on a mg/m² basis.

Use in Pregnancy

Category B3

There are insufficient clinical data to assess the possible effect of mirtazapine on pregnancy.

Oral dosing of pregnant rats with mirtazapine at 100 mg/kg/day was associated with a reduction in survival of the offspring, and an increased incidence of post-natal mortality. Mirtazapine was not teratogenic in rats at these dose levels, or in rabbits at oral doses up to 40 mg/kg/day

Although studies in animals have not shown any teratogenic effects of toxicological significance the safety of Remeron SolTab in human pregnancy has not been established. Epidemiological data have suggested that the use of SSRIs in pregnancy, particularly in late pregnancy, may increase the risk of persistent pulmonary hypertension in the newborn (PPHN). Although no studies have investigated the association of PPHN to mirtazapine treatment, this potential risk cannot be ruled out taking into account the related mechanism of action (increase in serotonin concentrations). Remeron SolTab should be used during pregnancy only if it is clearly needed. Women of child-bearing potential should employ an adequate method of contraception if taking Remeron SolTab.

Use in Lactation

Although animal experiments show that mirtazapine is excreted only in very small amounts in the milk, post natal mortality was increased when lactating rats were given mirtazapine orally at 100 mg/kg/day.

The use of Remeron SolTab in breast-feeding mothers is not recommended since no human data in breast milk are available.

INTERACTION WITH OTHER MEDICINES

Pharmacokinetic interactions

- Mirtazapine is extensively metabolised by CYP2D6 (resulting in the 8-hydroxy metabolite) and CYP3A4 (N-demethyl and N-oxide metabolites) and to a lesser extent by CYP1A2. An interaction study with healthy volunteers showed no influence of paroxetine, a CYP2D6 inhibitor, on mirtazapine pharmacokinetics in steady state.
- Coadministration of the potent inhibitor of CYP3A4, ketoconazole, in healthy male volunteers increased mirtazapine peak plasma concentration levels and AUC by approximately 40% and 50% respectively.
- When cimetidine (weak inhibitor of CYP1A2, CYP2D6 and CYP3A4) is administered with mirtazapine, the mean plasma concentration of mirtazapine may increase more than 50%. The mirtazapine dose may have to be decreased when concomitant treatment with cimetidine is started or increased when cimetidine treatment is ended. Caution should be exercised and the dose may have to be decreased when co-administering mirtazapine with potent CYP3A4 inhibitors, HIV protease inhibitors, azole antifungals, ketoconazole, erythromycin, cimetidine or nefazodone.
- Carbamazepine and phenytoin, inducers of CYP3A4, increased mirtazapine clearance about twofold, resulting in a decrease in mirtazapine plasma levels of 45-60%. When carbamazepine, phenytoin or another inducer of drug metabolism (such as rifampicin) is added to mirtazapine therapy, the mirtazapine dose may have to be increased. If treatment with an inducer is stopped, mirtazapine dose may have to be decreased.
- In in-vivo interaction studies, mirtazapine did not influence the pharmacokinetics of paroxetine (CYP2D6 substrates), carbamazepine or phenytoin (CYP3A4 inducers), amitriptyline, or cimetidine.

- In a mirtazapine and lithium interaction study, the steady state pharmacokinetics of lithium was not affected by coadministration of a single oral dose of 30mg of mirtazapine. Correspondingly, the single dose pharmacokinetics of mirtazapine was not affected by the lithium steady state.

Pharmacodynamic interactions

- Mirtazapine should not be administered concomitantly with MAO inhibitors or within two weeks after discontinuation of MAO inhibitor therapy. In the opposite way about two weeks should pass before patients treated with mirtazapine should be treated with MAO inhibitors (see **CONTRAINDICATIONS**). In addition, as with SSRIs, co-administration with other serotonergic active substances (L-tryptophan, triptans, tramadol, linezolid, methylene blue, SSRIs, venlafaxine, lithium and St. John's Wort – *Hypericum perforatum* – preparations) may lead to an incidence of serotonin associated effects (see **PRECAUTIONS**). Caution should be advised and a closer clinical monitoring is required when these active substances are combined with mirtazapine.
- Mirtazapine may potentiate the sedative effects of benzodiazepines and other sedatives (especially antipsychotics, antihistamine H1 antagonists, opioids). Caution should be taken when these drugs are prescribed together with mirtazapine.
- Mirtazapine may potentiate the central nervous dampening action of alcohol; patients using mirtazapine should therefore be advised to avoid alcohol during tasks which require concentration and alertness.
- Mirtazapine dosed at 30 mg daily caused a small but statistically significant increase of the INR in subjects treated with warfarin. Both at continuing stable doses and higher doses of mirtazapine, a more pronounced effect cannot be excluded. It is advisable to monitor the prothrombin time more carefully in case of concomitant treatment of warfarin with mirtazapine.
- The risk of QT prolongation and/or ventricular arrhythmias (e.g. Torsades de Pointes) may be increased with concomitant use of medicines which prolong the QTc interval (e.g. some antipsychotics and antibiotics) and in case of mirtazapine overdose.

From post-marketing experience it appears that serotonin syndrome occurs very rarely in patients treated with mirtazapine in combination with SSRIs or venlafaxine. If the combination is considered therapeutically necessary, dosage changes should be made with caution and there should be adequate close monitoring for early signs of serotonergic overstimulation.

Effects on ability to drive and use machines

Remeron SolTab may impair concentration and alertness (more commonly in the initial phase of treatment). Patients treated with Remeron SolTab should be cautioned about operating hazardous machinery, including automobiles, until they are reasonably certain that the treatment does not affect them adversely.

ADVERSE EFFECTS

Clinical Trials

Depressed patients display a number of symptoms that are associated with the illness itself. It is therefore sometimes difficult to ascertain which symptoms are a result of the illness itself and which are a result of treatment with mirtazapine.

System Organ Class	Common (>1/100 to <1/10)	Uncommon (>1/1,000 to ≤ 1/100)	Rare (>1/10,000 to ≤1/1,000)
Blood and the lymphatic system disorders			<ul style="list-style-type: none"> • Granulocytopenia • Agranulocytosis
Metabolism and nutrition disorders	<ul style="list-style-type: none"> • Increase in appetite 		
Psychiatric disorders			<ul style="list-style-type: none"> • Mania • Nightmares/ vivid dreams (paroniria)
Nervous system disorders	<ul style="list-style-type: none"> • Drowsiness/ sedation generally occurring during the first weeks 	<ul style="list-style-type: none"> • Dizziness • Headache 	<ul style="list-style-type: none"> • Epileptic seizures • Tremor • Convulsions (insults) • Myoclonus • Paraesthesia • Restless legs (hyperkinesia)
Vascular disorders			<ul style="list-style-type: none"> • Orthostatic hypotension
Hepatobiliary disorders		<ul style="list-style-type: none"> • Elevations in serum transaminase activities 	
Skin and subcutaneous tissue disorders			<ul style="list-style-type: none"> • Exanthema
Musculoskeletal connective tissue and bone disorders			<ul style="list-style-type: none"> • Arthralgia • Myalgia
General disorders and administration site conditions	<ul style="list-style-type: none"> • Generalised or local oedema 		
Investigations	<ul style="list-style-type: none"> • Weight gain 		

Post-marketing reports

Skin and subcutaneous tissue disorders: Stevens-Johnson syndrome, dermatitis bullous, erythema multiforme, toxic epidermal necrolysis, rash (including erythematous and maculopapular), rare cases of increased sweating, alopecia, pruritus and urticaria.

Musculoskeletal connective tissue and bone disorders: back pain, arthralgia, myalgia, rhabdomyolysis.

Nervous system disorders: lethargy, dysarthria, serotonin syndrome, somnolence (i.e drowsiness sedation), impaired concentration, dizziness, paraesthesia, headache, hyperkinesia, rare cases of; cerebrovascular disorder, convulsions, tremor and myoclonus, movement disorders.** Very rare cases of oral paraesthesia.

Psychiatric disorders: suicidal ideation***, suicidal behaviour***, confusion, agitation, aggression, paroniria, less common or rare occurrences of; nightmares/vivid dreams, hallucination, mania, depression, anxiety*, insomnia* and **psychomotor restlessness.

Gastrointestinal disorders: constipation, vomiting, pancreatitis, increased salivation nausea, diarrhoea, dry mouth, less common or rare cases of stomatitis, very rare cases of oral hypoaesthesia and mouth oedema.

Hepatobiliary disorders: hepatic function abnormal, elevated hepatic enzymes or transaminases, rare cases of jaundice, hepatitis.

Metabolism and nutrition disorders: hyponatraemia, increased appetite, rare cases of hypercholesterolaemia, hyperlipidaemia.

Cardiac disorders: tachycardia, palpitations, rare; arrhythmia, myocardial infarction, chest pain.

Vascular disorders: hypotension, dependent oedema, hypertension, orthostatic hypotension, rare cases of thromboembolic disorder, pulmonary embolism.

Blood and lymphatic system disorders: leukopenia, granulocytopenia, rare cases of agranulocytosis, (refer to **PRECAUTIONS**), rare cases of; thrombocytopenia, pancytopenia, anaemia, aplastic anaemia, eosinophilia and coagulation disorder.

Renal and urinary disorders: rare cases of urinary retention.

General disorders and administration site conditions: oedema including generalised, peripheral and face oedema; fatigue/asthenia, rare cases of pyrexia, syncope, chest pain and drug withdrawal symptoms.

Investigations: increases in gamma-glutamyltransferase levels, hypertriglyceridaemia, weight gain, increased creatine kinase.

Eye Disorders: Very rare cases of glaucoma.

*Upon treatment with antidepressants in general, anxiety and insomnia (which may be symptoms of depression) can develop or become aggravated. Under mirtazapine treatment, development or aggravation of anxiety and insomnia has been reported very rarely.

** Including akathisia, hyperkinesia.

***Cases of suicidal ideation and suicidal behaviours have been reported during mirtazapine therapy or early after treatment discontinuation (see **PRECAUTIONS**).

DOSAGE AND ADMINISTRATION

The tablets should be taken orally, if necessary with fluid, and swallowed without chewing.

Adults: Treatment should begin with 15 mg daily. The dosage generally needs to be increased to obtain an optimal clinical response. The effective daily dose is usually between 30 and 45 mg, but responses have been observed at 60 mg per day.

Elderly: The recommended dose is the same as that for adults. In elderly patients an increase in dosing should be done under close supervision to elicit a satisfactory and safe response.

The clearance of mirtazapine may be decreased in patients with renal or hepatic insufficiency. This should be taken into account when prescribing Remeron SolTab to this category of patients (refer to Pharmacokinetics).

Mirtazapine has a half-life of 20-40 hours and therefore Remeron SolTab is suitable for once-a-day administration. It should be taken preferably as a single night-time dose before going to bed. Remeron SolTab may also be given in sub-doses equally divided over the day (once in the morning and once at night-time).

Treatment should preferably be continued until the patient has been completely symptom-free for 4-6 months. After this, treatment can be gradually discontinued to avoid withdrawal symptoms (see **PRECAUTIONS**). Mirtazapine begins to exert its effect in general after 1-2 weeks of treatment. Treatment with an adequate dose should result in a positive response within 2-4 weeks. With an insufficient response, the dose can be increased up to the maximum dose. If there is no response within a further 2-4 weeks, then treatment should be stopped.

Children and adolescents (<18 years of age)

In placebo controlled trials, safety and efficacy of Remeron SolTab in the treatment of children and adolescents under the age of 18 years with major depressive disorder have not been established. Safety and efficacy in this population cannot be extrapolated from adult data. Therefore Remeron SolTab should not be used in children and adolescents under the age of 18 years.

OVERDOSAGE

Post-marketing experience concerning overdose with Remeron SolTab alone indicates that symptoms are usually mild. The symptoms of overdose are an exaggeration of the pharmacological action of mirtazapine and may include symptoms such as dizziness, impaired consciousness (confusion, disorientation, stupor, coma), agitation, tremor, tachycardia and hyper- and hypotension. As with all overdose attempts, the possibility of multiple drug ingestion should be borne in mind. As with antidepressants in general, serious outcomes, including fatalities, are possible at dosages much higher than the therapeutic dose, especially with mixed overdoses. In these cases QT prolongation and Torsade de Pointes have also been reported.

Overdose management

Cases of overdose should receive appropriate symptomatic and supportive therapy for vital functions. ECG monitoring should be undertaken.

PRESENTATION AND STORAGE CONDITIONS

Remeron SolTab tablets are round, white, flat faced bevelled-edge tablets marked with a code on one side, with a characteristic orange odour.

Remeron SolTab tablets are packed in a child-resistant, peel-to-open, rigid perforated blister, formed from a laminate of aluminium foil and plastic films sealed to a paper-based laminate of aluminium foil coated with a heat seal lacquer. Tablet pockets in the strip are separated by perforations.

The following packages are available:

15 mg (code $\begin{matrix} \text{TZ} \\ 1 \end{matrix}$) 12, 30, 48 and 90 tablets in peel-to-open blisters

30 mg (code $\begin{matrix} \text{TZ} \\ 2 \end{matrix}$) 12, 30, 48 and 90 tablets in peel-to-open blisters

45 mg (code $\begin{matrix} \text{TZ} \\ 4 \end{matrix}$) 12, 30, 48 and 90 tablets in peel-to-open blisters

Remeron SolTab should be stored in the dark and dry below 30°C.

NAME AND ADDRESS OF THE SPONSOR

Merck Sharp & Dohme (Australia) Pty Limited
Level 1, Building A, 26 Talavera Road
Macquarie Park, NSW 2113
Australia

POISON SCHEDULE OF THE MEDICINE

Schedule 4 – Prescription Only Medicine

DATE OF FIRST INCLUSION IN THE ARTG

09 August 2002

DATE OF MOST RECENT AMENDMENT

27 May 2016