

AUSTRALIAN PRODUCT INFORMATION

SINGULAIR®

(MONTELUKAST SODIUM) Tablets and Granules

1 NAME OF THE MEDICINE

Montelukast sodium

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each 10 mg film-coated tablet contains 10.4 mg montelukast sodium, which is equivalent to 10.0 mg of montelukast, the free acid.

Excipients with known effects:

- lactose monohydrate

Each 5 mg chewable tablet contains 5.2 mg montelukast sodium, which is equivalent to 5.0 mg of montelukast, the free acid.

Each 4 mg chewable tablet contains 4.16 mg of montelukast sodium, which is equivalent to 4.0 mg of montelukast, the free acid.

- Excipients with known effects:
- aspartame

Each packet of 4 mg oral granules contains 4.16 mg of montelukast sodium, which is equivalent to 4.0 mg of montelukast, the free acid.

For the full list of excipients, see **Section 6.1 List of Excipients**.

3 PHARMACEUTICAL FORM

10 mg film-coated tablet – A beige biconvex rounded square film-coated tablet engraved with SINGULAIR on one side and MSD 117 on the other.

5 mg chewable tablet - A pink biconvex round chewable tablet engraved with SINGULAIR on one side and MSD 275 on the other.

4 mg chewable tablet - A pink, oval, biconvex chewable tablet with SINGULAIR debossed on one side and MSD 711 on the other.

4 mg oral granules - Coarse, white free-flowing granules sealed in a foil packet.

4 CLINICAL PARTICULARS

4.1 THERAPEUTIC INDICATIONS

Prophylaxis and treatment of chronic asthma in adults and children 2 years of age and older.

Symptomatic treatment of seasonal allergic rhinitis.

4.2 DOSE AND METHOD OF ADMINISTRATION

Asthma and/or Seasonal Allergic Rhinitis

SINGULAIR should be taken once daily.

For asthma, the dose should be taken in the evening.

For seasonal allergic rhinitis, the time of administration may be individualised to suit patient needs.

Patients with both asthma and seasonal allergic rhinitis should take only one tablet daily in the evening.

Adults 15 Years of Age and Older

The dosage for adults 15 years of age and older is one 10 mg tablet daily.

Paediatric Patients 6 to 14 Years of Age

The dosage for paediatric patients 6 to 14 years of age is one 5 mg chewable tablet daily. Chew the tablets. Do not swallow whole.

Paediatric Patients 2 to 5 Years of Age

The dosage for paediatric patients 2 to 5 years of age is one 4 mg chewable tablet daily or one packet of 4 mg oral granules daily. Chew the tablets. Do not swallow whole.

Administration of Oral Granules

SINGULAIR oral granules can be administered either directly in the mouth, or mixed with a spoonful of cold or room temperature soft food (e.g. applesauce). The packet should not be opened until ready to use. After opening the packet, the full dose of SINGULAIR oral granules must be administered immediately (within 15 minutes). If mixed with food, SINGULAIR oral granules must not be stored for future use. SINGULAIR oral granules are not intended to be dissolved in liquid for administration. However, liquids may be taken subsequent to administration.

General Recommendations

The therapeutic effect of SINGULAIR on parameters of asthma control occurs within one day. SINGULAIR tablets, chewable tablets and oral granules can be taken with or without food. Patients should be advised to continue taking SINGULAIR daily when their asthma is controlled, as well as during periods of worsening asthma.

No dosage adjustment is necessary for paediatric patients, for the elderly, for patients with renal insufficiency, or mild to moderate hepatic impairment, or for patients of either gender.

Therapy with SINGULAIR in Relation to Other Treatments for Asthma

SINGULAIR can be added to a patient's existing treatment regimen.

Reduction in Concomitant Therapy:

Bronchodilator Treatments: SINGULAIR can be added to the treatment regimen of patients who are not adequately controlled on bronchodilator alone. When a clinical response is evident (usually after the first dose), the patient's bronchodilator therapy can be reduced as tolerated.

Inhaled Corticosteroids: Treatment with SINGULAIR provides additional clinical benefit to patients treated with inhaled corticosteroids. A reduction in the corticosteroid dose can be made as tolerated. The dose should be reduced gradually with medical supervision. In some patients, the dose of inhaled corticosteroids can be tapered off completely. SINGULAIR should not be abruptly substituted for inhaled corticosteroids.

4.3 CONTRAINDICATIONS

Hypersensitivity to any component of these products.

4.4 SPECIAL WARNINGS AND PRECAUTIONS FOR USE

The efficacy of oral SINGULAIR for the treatment of acute asthma attacks has not been established. Therefore, oral tablets of SINGULAIR should not be relied upon to treat acute asthma attacks. Patients should be advised to have appropriate rescue medication available.

While the dose of concomitant inhaled corticosteroid may be reduced gradually under medical supervision, SINGULAIR should not be abruptly substituted for inhaled or oral corticosteroids (see **Section 5.1 Pharmacodynamic Properties, Clinical Trials - Asthma**).

Neuropsychiatric events

Neuropsychiatric events have been reported in adult, adolescent, and paediatric patients taking SINGULAIR. Post-marketing reports with SINGULAIR use include agitation, aggressive behaviour or hostility, anxiousness, depression, dream abnormalities, hallucinations, insomnia, irritability, restlessness, somnambulism, suicidal thinking and behaviour (including suicidality), and tremor. The clinical details of some post-marketing reports involving SINGULAIR appear consistent with a drug-induced effect.

Patients and prescribers should be alert for neuropsychiatric events. Patients should be instructed to notify their prescriber if these changes occur. Prescribers should carefully evaluate the risks and benefits of continuing treatment with SINGULAIR if such events occur.

Eosinophilic conditions

In rare cases patients receiving anti-asthma agents including leukotriene receptor antagonists, have experienced one or more of the following: eosinophilia, vasculitic rash, worsening pulmonary symptoms, cardiac complications, and/or neuropathy sometimes diagnosed as Churg-Strauss syndrome, a systemic eosinophilic vasculitis. These cases have been sometimes associated with the reduction or withdrawal of oral corticosteroid therapy. Although a causal relationship with leukotriene receptor antagonism has not been established, caution and appropriate clinical monitoring are recommended in patients receiving SINGULAIR.

4 mg and 5 mg chewable tablets contain aspartame which is a source of phenylalanine (0.842 mg phenylalanine per 5 mg chewable tablet, and 0.674 mg phenylalanine per 4 mg chewable tablet).

Although montelukast is effective in improving airway function in asthmatics with documented aspirin sensitivity, it has not been shown to modify bronchoconstrictor response to aspirin challenge and other non-steroidal anti-inflammatory drugs in aspirin-sensitive asthmatic patients. Therefore, patients with known aspirin sensitivity should continue avoidance of aspirin or non-steroidal anti-inflammatory agents while taking SINGULAIR. (**See Section 5.1 Pharmacodynamic Properties, Clinical studies - Asthma**).

Use in hepatic impairment

No dosage adjustment is necessary for the elderly or for patients with mild to moderate hepatic insufficiency. There are no clinical data in patients with severe hepatic insufficiency (Child-Pugh score > 9).

Use in renal impairment

Because montelukast and its metabolites are eliminated by the biliary route, no dose adjustment is anticipated to be necessary in patients with renal impairment. Studies in patients with renal impairment have not been undertaken.

Use in the elderly

In clinical studies, there were no age-related differences in the efficacy or safety profiles of SINGULAIR.

Paediatric use

In asthma: SINGULAIR has been studied in paediatric patients six months to 14 years of age (see **Section 4.2 Dose and Method of Administration**). Safety and effectiveness in paediatric patients younger than six months of age have not been studied. In studies investigating the effect of SINGULAIR on the growth rate of paediatric patients, it has been shown in one study that SINGULAIR does not affect the growth rate of paediatric patients when given for up to 56 weeks. The long term clinical relevance of the growth rates studies is unknown.

In seasonal allergic rhinitis: SINGULAIR has been studied in paediatric patients 2 to 14 years of age (see **Section 4.2 Dose and Method of Administration**). Safety in paediatric patients younger than two years of age has not been studied.

Effects on laboratory tests

No data available.

4.5 INTERACTIONS WITH OTHER MEDICINES AND OTHER FORMS OF INTERACTIONS

Relatively high concentrations of montelukast competitively inhibit the activity of cytochromes P450 3A4 and 2C9. However, these concentrations are at least 15 fold higher than the peak plasma concentrations attained following a 10 mg oral dose of montelukast. Based on these and other *in vitro* results in human liver microsomes, therapeutic plasma concentrations of montelukast should not be expected to inhibit cytochromes P450 3A4, 2C9, 1A2, 2A6, 2C19 or 2D6. Theophylline plasma concentration was not affected by the recommended dose of SINGULAIR (10 mg once daily). At 20 and 60 fold above the recommended dose, plasma concentration of concomitant theophylline was decreased. Theophylline dose adjustment or a change in the frequency of plasma theophylline monitoring is not necessary at the recommended dose of SINGULAIR.

SINGULAIR may be administered with other therapies routinely used in the prophylaxis and chronic treatment of asthma, and in the treatment of allergic rhinitis. In drug-interaction studies, the recommended clinical dose of montelukast did not have clinically important effects on the pharmacokinetics of the following drugs: theophylline, prednisone, prednisolone, oral contraceptives (ethinyl estradiol/ norethisterone 35/1), terfenadine, digoxin and warfarin. The effects of concomitant administration of montelukast and macrolide antimicrobials have not been studied.

The area under the plasma concentration-time curve (AUC) for montelukast was decreased approximately 40% in subjects with co-administration of phenobarbital. No dosage adjustment for SINGULAIR is recommended.

In vitro studies have shown that montelukast is an inhibitor of CYP 2C8. However, data from a clinical drug-drug interaction study involving montelukast and rosiglitazone (a probe substrate representative of drugs primarily metabolized by CYP2C8) demonstrated that montelukast does not inhibit CYP2C8 *in vivo*. Therefore, montelukast is not anticipated to alter

the metabolism of drugs metabolized by this enzyme (e.g., paclitaxel, rosiglitazone, repaglinide).

In vitro studies have shown that montelukast is a substrate of CYP 2C8, 2C9, and 3A4. Data from a clinical drug-drug interaction study involving montelukast and gemfibrozil (an inhibitor of both CYP 2C8 and 2C9) demonstrated that gemfibrozil increased the systemic exposure of montelukast by 4.4-fold. Co-administration of itraconazole, a strong CYP 3A4 inhibitor, with gemfibrozil and montelukast did not further increase the systemic exposure of montelukast. Based on available clinical experience no dosage adjustment of montelukast is required upon co-administration with gemfibrozil or other potent inhibitors of CYP 2C8. Specific *in vivo* data are not available for other known inhibitors of CYP 2C8 (e.g. trimethoprim). Although clinically important drug interactions are not anticipated based on these findings, systemic exposure to montelukast is potentially increased and the physician should be aware of a possible increase in adverse reactions (see **Section 4.9 Overdose**). Co-administration of montelukast with itraconazole alone resulted in no significant increase in the systemic exposure of montelukast.

Although additional specific interaction studies were not performed, SINGULAIR was used concomitantly with a wide range of commonly prescribed drugs in clinical studies without evidence of clinical adverse interactions. These medications included thyroid hormones, sedative hypnotics, nonsteroidal anti-inflammatory agents, benzodiazepines and decongestants.

4.6 FERTILITY, PREGNANCY AND LACTATION

Effects on fertility

Fertility and reproductive performance were not affected in studies with male rats given oral doses of up to 800 mg/kg/day, but fecundity was slightly reduced in female rats doses orally at 200 mg/kg/day. The no-effect dose for the latter effect was 100 mg/kg/day, corresponding to systemic exposure, in terms of plasma AUC for parent drug, at least 20 times higher than that in women at recommended dose levels.

Use in pregnancy

(Category B1)

In animal studies, montelukast sodium had no adverse effects on embryofoetal development at oral doses up to 400 mg/kg/day in rats or up to 100 mg/kg/day in rabbits. Retardation of foetal growth and development was observed in rabbits dosed at 200 mg/kg/day, a dose level associated with severe maternal toxicity. Foetal exposure of montelukast was demonstrated in both species. SINGULAIR has not been studied in pregnant women. SINGULAIR should be used during pregnancy only if clearly needed.

During worldwide marketing experience, congenital limb defects have been rarely reported in the offspring of women being treated with SINGULAIR during pregnancy. Most of these women were also taking other asthma medications during their pregnancy. A causal relationship between these events and SINGULAIR has not been established.

Use in lactation

Studies in lactating rats have shown that montelukast is excreted into milk following oral doses of 100 and 200 mg/kg/day, and growth of the pups was slightly inhibited at the higher dose level. It is not known if SINGULAIR is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when SINGULAIR is given to a nursing mother.

4.7 EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

There is no evidence that SINGULAIR affects the ability to drive and use machines. However, adverse effects of these medicines include dizziness and drowsiness, which could affect the ability to drive or use machines (see **Section 4.8 ADVERSE EFFECT (UNDESIRABLE EFFECTS)**).

4.8 ADVERSE EFFECTS (UNDESIRABLE EFFECTS)

SINGULAIR has been generally well tolerated. Side effects, which usually were mild, generally did not require discontinuation of therapy. The overall incidence of side effects reported with SINGULAIR was comparable to placebo.

Adults 15 years of Age and Older with Asthma

SINGULAIR has been evaluated for safety in approximately 2600 adult patients 15 years of age and older in clinical studies. In two similarly designed, 12 week placebo-controlled clinical studies, the only adverse experiences reported as drug-related in $\geq 1\%$ of patients treated with SINGULAIR and at a greater incidence than in patients treated with placebo were abdominal pain and headache. The incidences of these events were not significantly different in the two treatment groups.

In placebo-controlled clinical studies, the following adverse experiences reported with SINGULAIR occurred in $\geq 1\%$ of patients and at an incidence greater than or equal to that in patients treated with placebo, regardless of drug relationship:

Table 1
ADVERSE EXPERIENCES OCCURRING IN $\geq 1\%$ OF PATIENTS WITH AN INCIDENCE
GREATER THAN OR EQUAL TO THAT IN PATIENTS TREATED WITH PLACEBO,
REGARDLESS OF DRUG RELATIONSHIP

	SINGULAIR 10 mg/day (%) (n = 1955)	Placebo (%) (n = 1180)
Body As A Whole		
Asthenia/fatigue	1.8	1.2
Fever	1.5	0.9
Pain, abdominal	2.9	2.5
Trauma	1.0	0.8
Digestive System Disorders		
Diarrhoea	3.1	3.1
Dyspepsia	2.1	1.1
Gastroenteritis, infectious	1.5	0.5
Pain, dental	1.7	1.0
Nervous System/Psychiatric		
Dizziness	1.9	1.4
Headache	18.4	18.1
Insomnia	1.3	1.3
Respiratory System Disorders		
Congestion, nasal	1.6	1.3
Cough	2.7	2.4
Influenza	4.2	3.9
Skin/Skin Appendages Disorder		
Rash	1.6	1.2

Laboratory Adverse Experiences*		
ALT increased	2.1	2.0
AST increased	1.6	1.2
Pyuria	1.0	0.9

* Number of patients tested (SINGULAIR and placebo, respectively): ALT and AST, 1935, 1170; pyuria, 1924, 1159.

Cumulatively, 544 patients were treated with SINGULAIR for at least 6 months, 253 for one year and 21 for two years in clinical studies. With prolonged treatment, the adverse experience profile did not change.

Paediatric Patients 6 to 14 Years of Age with Asthma

SINGULAIR has been evaluated for safety in approximately 970 paediatric patients 6 to 14 years of age. The safety profile in paediatric patients is generally similar to the adult safety profile and to placebo. Cumulatively, 263 paediatric patients 6-14 years of age were treated with SINGULAIR for at least 3 months, 164 for 6 months or longer in clinical studies. The safety profile in paediatric patients is generally similar to the adult safety profile and to placebo. With prolonged treatment, the adverse experience profile did not change.

In a 56 week active-controlled study comparing montelukast to inhaled fluticasone in paediatric patients 6-14 years of age with mild persistent asthma, the safety profile was consistent with the safety profile previously described for SINGULAIR. In the study, the number of patients with asthma symptoms after treatment was 166 (33.5%) patients in the montelukast treatment group and 135 (27.1%) patients in the fluticasone treatment group.

In studies evaluating growth rate, the safety profile in these paediatric patients was consistent with the safety profile previously described for SINGULAIR.

Paediatric Patients 2 to 5 Years of Age with Asthma

SINGULAIR has been evaluated in 573 paediatric patients 2 to 5 years of age. In a 12-week placebo-controlled clinical study, the only adverse experience reported as drug related in > 1% of patients treated with SINGULAIR and at a greater incidence than in patients treated with placebo was thirst. The incidence of thirst was not significantly different in the two treatment groups.

Cumulatively, 502 paediatric patients 2 to 5 years of age were treated with SINGULAIR for at least 3 months, 338 for 6 months or longer, and 256 patients for 12 months or longer. With prolonged treatment, the adverse experience profile did not change.

Adults 15 Years of Age and Older with Seasonal Allergic Rhinitis

SINGULAIR has been evaluated in 2199 adult patients 15 years of age and older for the treatment of seasonal allergic rhinitis in clinical studies. SINGULAIR administered once daily in the morning or in the evening was generally well tolerated with a safety profile similar to that of placebo. In placebo-controlled clinical studies, no adverse experiences reported as drug related in $\geq 1\%$ of patients treated with SINGULAIR and at a greater incidence than in patients treated with placebo were observed. In a 4-week, placebo-controlled clinical study, the safety profile was consistent with that observed in 2-week studies. The incidence of somnolence was similar to that of placebo in all studies.

Paediatric Patients 2 to 14 Years of Age with Seasonal Allergic Rhinitis

SINGULAIR has been evaluated in 280 paediatric patients 2 to 14 years of age for the treatment of seasonal allergic rhinitis in a 2-week, placebo-controlled, clinical study. SINGULAIR administered once daily in the evening was generally well tolerated with a safety

profile similar to that of placebo. In this study, no adverse experiences reported as drug related in $\geq 1\%$ of patients with SINGULAIR and at a greater incidence than in patients treated with placebo were observed.

Adults 15 Years of Age and Older with Asthma and Seasonal Allergic Rhinitis

SINGULAIR 10 mg film-coated tablets have been evaluated in approximately 400 asthmatic patients 15 years of age and older with seasonal allergic rhinitis. The safety profile in asthmatic patients with seasonal allergic rhinitis was consistent with that observed in patients with asthma.

Post-Marketing Experience

The following additional side effects have been reported in post-marketing use:

Infections and infestations: upper respiratory tract infection

Blood and lymphatic system disorders: increased bleeding tendency, thrombocytopenia

Immune system disorders: hypersensitivity reactions including anaphylaxis, very rarely hepatic eosinophilic infiltration

Psychiatric disorders: agitation including aggressive behaviour or hostility, anxiousness, depression, disorientation, disturbance in attention, dream abnormalities, hallucinations, insomnia, memory impairment, obsessive-compulsive symptoms, psychomotor hyperactivity (including irritability, restlessness, and tremor), somnambulism (sleep walking), suicidal thinking and behaviour (suicidality), tic

Nervous system disorders: dizziness, drowsiness, paraesthesia/hypoesthesia, very rarely seizure

Cardiac disorders: palpitations

Respiratory, thoracic and mediastinal disorders: epistaxis, pulmonary eosinophilia

Gastrointestinal disorders: diarrhoea, dyspepsia, nausea, vomiting

Hepatobiliary disorders: increased ALT and AST, very rarely hepatitis (including cholestatic, hepatocellular, and mixed-pattern liver injury)

Skin and subcutaneous tissue disorders: angioedema, bruising, erythema multiforme[†], erythema nodosum, pruritus, rash, urticaria

Musculoskeletal and connective tissue disorders: arthralgia, myalgia including muscle cramps

Renal and urinary disorders: enuresis in children

General disorders and administration site conditions: oedema, pyrexia

In rare cases, patients on therapy with SINGULAIR may present with systemic eosinophilia, sometimes presenting with clinical features of vasculitis consistent with Churg-Strauss syndrome. These events usually, but not always, have been associated with the reduction of oral corticosteroid therapy. A causal association between SINGULAIR and these underlying conditions has not been established (see **Section 4.4 Special Warnings and Precautions for Use**).

Reporting suspected adverse effects

Reporting suspected adverse reactions after registration of the medicinal product is important. It allows continued monitoring of the benefit-risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions at www.tga.gov.au/reporting-problems.

4.9 OVERDOSE

No specific information is available on the treatment of overdose with SINGULAIR. In chronic asthma studies, SINGULAIR has been administered at doses up to 200 mg/day to adult patients for 22 weeks and in short-term studies, up to 900 mg/day to patients for approximately one week without clinically important adverse experiences.

There have been reports of acute overdose in postmarketing experience and clinical studies with SINGULAIR. These include reports in adults and children with a dose as high as 1000 mg. The clinical and laboratory findings observed were consistent with the safety profile in adults and paediatric patients.

There were no adverse experiences in the majority of overdose reports.

The most frequently occurring adverse experiences were consistent with the safety profile of SINGULAIR and included abdominal pain, somnolence, thirst, headache, vomiting, and psychomotor hyperactivity.

It is not known whether montelukast is dialyzable by peritoneal- or hemodialysis.

For information on the management of overdose, contact the Poisons Information Centre on 131126 (Australia).

5 PHARMACOLOGICAL PROPERTIES

5.1 PHARMACODYNAMIC PROPERTIES

Mechanism of action

The cysteinyl leukotrienes (LTC₄, LTD₄, LTE₄), are potent inflammatory eicosanoids released from various cells including mast cells and eosinophils. These important pro-asthmatic mediators bind to cysteinyl leukotriene (CysLT) receptors. The CysLT type-1 (CysLT₁) receptor is found in the human airway (including airway smooth muscle cells and airway macrophages) and on other pro-inflammatory cells (including eosinophils and certain myeloid stem cells). CysLTs have been correlated with the pathophysiology of asthma and allergic rhinitis. In asthma, leukotriene-mediated effects include a number of airway actions, including bronchoconstriction, mucous secretion, vascular permeability, and eosinophil recruitment. In allergic rhinitis, CysLTs are released from the nasal mucosa after allergen exposure during both early- and late-phase reactions and are associated with symptoms of allergic rhinitis. Intranasal challenge with CysLTs has been shown to increase nasal airway resistance and symptoms of nasal obstruction. The clinical relevance of intranasal challenge studies is unknown.

Montelukast is an orally active compound which has been shown in asthmatic patients to reduce peripheral blood eosinophil counts and sputum eosinophils, which are parameters of asthmatic inflammation. The effect of montelukast on reduction of peripheral blood eosinophils was comparable to inhaled corticosteroids. Based on biochemical and pharmacological bioassays, it binds with high affinity and selectivity to the CysLT₁ receptor (in preference to other pharmacologically important airway receptors such as the prostanoid, cholinergic, or β -adrenergic receptor). Montelukast potently inhibits physiologic actions of LTC₄, LTD₄, and LTE₄ at the CysLT₁ receptor without any agonist activity.

In asthmatic patients, montelukast causes potent inhibition of airway cysteinyl leukotriene receptors as demonstrated by the ability to inhibit bronchoconstriction due to inhaled LTD₄. A dose of 5 mg causes substantial blockage of LTD₄-induced bronchoconstriction. Montelukast causes bronchodilation within 2 hours of oral administration. β -agonists caused additive effects when added to montelukast.

Clinical trials

Clinical Studies - Asthma

In clinical studies, SINGULAIR is effective in adult and paediatric patients for the prophylaxis and chronic treatment of asthma, including protection against day- and nighttime symptoms, the treatment of aspirin-sensitive asthmatic patients, and the prevention of exercise-induced bronchoconstriction. SINGULAIR is effective alone or in combination with other prophylactic agents used in the maintenance treatment of asthma. SINGULAIR and inhaled corticosteroid may be used concomitantly with additive effects to control asthma or to reduce the inhaled corticosteroid dose while maintaining clinical stability. SINGULAIR is a preventative agent which should be used in addition to other drugs for the management of asthma.

Adults 15 years of age and older

In two similarly-designed 12-week double-blind placebo-controlled studies in adults asthmatic patients 15 years of age and older, SINGULAIR, 10 mg once daily in the evening, demonstrated significant improvements in parameters of asthma control measuring asthma symptoms, asthma-related outcomes, respiratory function and "as-needed" β -agonist use.

SINGULAIR significantly improved patient-reported daytime symptoms and nocturnal awakenings, compared with placebo. Asthma-specific outcomes, including asthma attacks, corticosteroid rescue, discontinuations due to worsening asthma, asthma exacerbations and asthma-free days were also significantly better than placebo.

Physicians' and patients' global asthma evaluations and asthma-specific quality-of-life evaluations (in all domains, including normal daily activity and asthma symptoms) were significantly better than placebo.

SINGULAIR caused significant improvements in morning forced expiratory volume in 1 second (FEV₁), AM and PM peak expiratory flow rate (PEFR) and significantly decreased the use of 'as needed' β -agonist, compared with placebo.

A comparison of SINGULAIR and inhaled beclomethasone (200 μ g twice daily with a spacer device) demonstrated that SINGULAIR had a more rapid initial response (within the first day compared with 7-10 days for beclomethasone). While both treatments provided significantly and clinically important changes, the overall beclomethasone effect was larger over the 12 weeks duration of the study. The difference in response is, in part, a result of a small percentage of patients treated with beclomethasone (16.7%) that had a \geq 30% increase in FEV₁). However in a high proportion of patients the response was comparable, for example, 42% of patients taking montelukast compared with 50% of patients taking beclomethasone achieved an 11% or more increase in FEV₁.

The treatment effect was achieved after the first dose and was maintained throughout the 24 hour dosing interval. Treatment effect also remained constant during continuous once-daily administration in extension studies for up to one year. Withdrawal of SINGULAIR after 12-weeks of use did not cause rebound worsening of asthma.

Paediatric patients 6 to 14 years of age

In paediatric patients 6 to 14 years of age, one 5 mg chewable tablet daily in the evening, significantly decreased asthma exacerbations, and improved parents' global evaluations and

the paediatric asthma-specific quality-of-life evaluations, compared with placebo. SINGULAIR also significantly improved morning FEV₁ and decreased total daily “as-needed” β-agonist use. Treatment effect was achieved after the first dose and remained constant during once-daily administration for up to 6 months.

Growth Rate in Paediatric Patients

Two controlled clinical studies have demonstrated that montelukast did not affect the growth rate in paediatric patients with asthma. In a short term study of children aged 6 to 11 years, growth rate as measured by lower leg length growth was similar in patients treated with montelukast 5 mg once daily for 3 weeks compared with placebo, and was significantly lower in patients treated with inhaled budesonide (200 µg twice daily) for 3 weeks, compared with placebo. In a 56-week study in children aged 6 to 8 years, linear growth rate was similar in patients treated with montelukast 5 mg once daily and placebo (LS means for montelukast and placebo: 5.67 and 5.64 cm/year, respectively), and was significantly lower (LS mean: 4.86 cm/year) in patients treated with inhaled beclomethasone (200 µg twice daily), compared with placebo [difference in LS means (95% CI): -0.78 (-1.06, -0.49) cm/year]. The long term clinical relevance of these studies is unknown.

Paediatric patients 2 to 5 years of age

In a 12-week, placebo-controlled study in paediatric patients 2 to 5 years of age, SINGULAIR 4 mg once daily consistently improved parameters of asthma control irrespective of concomitant controller therapy use compared with placebo. Sixty percent of patients were not on any other controller therapy. SINGULAIR significantly improved daytime symptoms (including coughing, wheezing, trouble breathing and activity limitation) and nighttime symptoms compared with placebo (mean baseline daytime scores were SINGULAIR 0.98 and placebo 0.95 on a 0-5 scale; mean change from baseline: SINGULAIR -0.37 vs placebo -0.25 [p=0.003] and mean baseline nighttime scores were SINGULAIR 1.18 and placebo 1.20 on a 0-4 scale; mean change from baseline: SINGULAIR -0.41 vs placebo -0.30 [p=0.026]).

SINGULAIR also significantly decreased “as-needed” β-agonist use (percentage of days used, SINGULAIR 50.09 vs placebo 56.34 [p=0.001]) and corticosteroid rescue (percentage of patients with corticosteroid rescue, SINGULAIR 19.09 vs placebo 28.07 [p=0.008]) compared with placebo. Patients receiving SINGULAIR had significantly more days without asthma than those receiving placebo (percentage of days without, SINGULAIR 30.50 vs placebo 23.63 [p=0.002]). A treatment effect was achieved after the first dose. In addition, total blood eosinophil counts were significantly decreased (for total blood eosinophil counts, the between group difference in LS means was -0.04 10³/µL, p=0.034).

Effects in patients on concomitant inhaled corticosteroids

Separate studies in adults demonstrated the ability of SINGULAIR to add to the clinical effect of inhaled corticosteroid and allow steroid tapering when used concomitantly. In a placebo-controlled study, stable, asthmatic patients taking initial inhaled corticosteroid doses of approximately 1600 µg per day reduced their steroid use by approximately 37% during a placebo run-in period. SINGULAIR allowed a further 47% reduction of the inhaled corticosteroid dose, compared with 30% for placebo. In another study, SINGULAIR provided additional clinical benefit to a similar population of patients maintained but not adequately controlled on inhaled corticosteroid (beclomethasone 400 µg per day). Complete abrupt removal of beclomethasone in patients receiving both SINGULAIR and beclomethasone caused clinical deterioration in some patients, suggesting that tapering as tolerated rather than abrupt removal is preferred.

Aspirin-sensitive patients

In aspirin-sensitive patients, nearly all of whom were receiving concomitant inhaled and/or oral corticosteroids, SINGULAIR resulted in significant improvement in the parameters of asthma control.

The effect of SINGULAIR on the bronchoconstrictor response to aspirin challenge or other non-steroidal anti-inflammatory drugs in aspirin-sensitive asthmatic patients has not been evaluated (see **Section 4.4 Special Warnings and Precautions for Use**).

Effects on exercise-induced bronchoconstriction

SINGULAIR, 10 mg once daily, protected against exercise-induced bronchoconstriction (EIB) in adults 15 years of age and older. In a 12-week study, SINGULAIR significantly inhibited the extent and duration of fall in FEV₁ over 60 minutes after exercise, the maximal percent fall in FEV₁ after exercise, and the time to recovery to within 5% of the pre-exercise FEV₁. Protection was consistent through the treatment period indicating that tolerance did not occur. In a separate cross-over study, protection was observed after two once-daily doses.

In paediatric patients 6 to 14 years of age, using the 5 mg chewable tablet, a similarly designed cross-over study demonstrated similar protection and the protection was maintained throughout the dosing interval (24 hours).

Effects on antigen challenge and eosinophils

In clinical studies SINGULAIR inhibited both early- and late-phase bronchoconstriction due to antigen challenge. Because inflammatory cell (eosinophil) infiltration is an important feature of asthma, the effects of SINGULAIR on eosinophils in the peripheral blood and airway were examined. In Phase IIb/III clinical studies, SINGULAIR significantly decreased peripheral blood eosinophils approximately 15% from baseline, compared with placebo. In paediatric patients 6 to 14 years of age, SINGULAIR decreased peripheral blood eosinophils 13% over the 8-week treatment period, compared with placebo. SINGULAIR also significantly decreased airway eosinophils in sputum, compared with placebo. In this study, peripheral blood eosinophils decreased and clinical asthma endpoints improved with treatment with SINGULAIR.

Effects in Patients with Asthma and Seasonal Allergic Rhinitis

In a two week, placebo controlled, double blind, clinical study in adult asthmatic patients 15 years of age and older with concomitant seasonal allergic rhinitis, montelukast 10 mg tablets administered once daily demonstrated a statistically significant improvement in the primary composite variable, Daily Rhinitis Symptoms score (average of the Daytime Nasal Symptoms score [mean of nasal congestion, rhinorrhea, sneezing, nasal itching] and the Nighttime Symptoms score [mean of nasal congestion upon awakening, difficulty going to sleep, and nighttime awakenings scores]) (see TABLE 2). The mean difference in improvement between the two treatments was - 0.12 points (95% CI: - 0.18, -0.06) on a scale of 0.00 to 3.00. A change of 1.00 would correspond to an improvement of symptoms from moderate to mild, or from severe to moderate.

Table 2
Daily Rhinitis Symptom Score

Treatment Group	N	Change from Baseline Mean (SD)	Treatment Difference: Montelukast Minus Placebo	
			LS Mean Difference (95% CI)	p-Value

Montelukast	412	-0.35 (0.48)	-0.12 (-0.18, -0.06)	≤ 0.001
Placebo	413	-0.24 (0.46)		

Clinical Studies - Seasonal Allergic Rhinitis

Adults 15 years of age and older

The efficacy of SINGULAIR for the treatment of seasonal allergic rhinitis was investigated in seven similarly designed randomised, 2-week, double-blind, placebo-controlled trials including 4924 patients (1751 patients were treated with SINGULAIR). Patients were 15 years of age and older with a history of seasonal allergic rhinitis, a positive skin test to at least one relevant seasonal allergen, and active symptoms of seasonal allergic rhinitis at study initiation.

Two of the three pivotal studies showed a significant reduction in daytime nasal symptoms scores with SINGULAIR 10 mg tablets compared to placebo.

In a combined analysis of the three pivotal studies, SINGULAIR 10 mg tablets administered to 1189 patients once daily in the evening resulted in a statistically significant improvement in the primary endpoint, daytime nasal symptoms score, and its individual components (nasal congestion, rhinorrhea, nasal itching, and sneezing); nighttime symptoms score, and its individual components (nasal congestion upon awakening, difficulty going to sleep, and nighttime awakenings); daytime eye symptoms score, and its individual components (tearing, itchy, red, and puffy eyes); global evaluation of allergic rhinitis by patients and by physicians; and composite symptoms score (composed of the daytime nasal and nighttime symptoms scores), compared with placebo.

The efficacy results of one trial are shown below. The mean changes from baseline in daytime nasal symptoms score in the treatment groups that received SINGULAIR tablets, loratadine and placebo are shown in Table 3. The mean change from baseline for SINGULAIR was 50% larger than the mean change from baseline for placebo.

Table 3
Effects of SINGULAIR on Daytime Nasal Symptoms Score* in a Placebo- and Active-controlled Trial in Patients with Seasonal Allergic Rhinitis

Treatment Group (N)	Baseline Mean Score	Mean Change from Baseline	Difference Between Treatment and Placebo (95% CI) Least-Squares Mean
SINGULAIR 10 mg (344)	2.09	-0.39	-0.13 [‡] (-0.21, -0.06)
Placebo (351)	2.10	-0.26	N.A.
Active Control [†] (Loratadine 10 mg) (599)	2.06	-0.46	-0.24 [‡] (-0.31, -0.17)

* Average of individual scores of nasal congestion, rhinorrhea, nasal itching, sneezing as assessed by patients on a 0-3 categorical scale.

† The study was not designed for statistical comparison between SINGULAIR and the active control (loratadine).

‡ Statistically different from placebo (p≤0.001).

The efficacy of SINGULAIR in the treatment of seasonal allergic rhinitis was investigated in a separate 4-week study in which the primary objective of the study was to determine the treatment effect of SINGULAIR 10 mg administered once daily in the morning, compared to placebo, in patients with seasonal allergic rhinitis over the first 2 weeks of treatment. The

efficacy of SINGULAIR over the initial 2 weeks was significantly different from placebo and consistent with the effect observed in studies using evening dosing (see Table 4). Additionally, the effect over the entire 4 weeks was consistent with the 2-week results (see Figure 1).

Table 4
Effects of SINGULAIR on Daytime Nasal Symptoms Score* in a Placebo- and Active-controlled Trial in Patients with Seasonal Allergic Rhinitis


Treatment Group (N)	Baseline Mean Score	Mean Change from Baseline	Difference Between Treatment and Placebo (95% CI) Least-Squares Mean
SINGULAIR 10 mg (445)	2.19	-0.32	-0.10 ^{††} (-0.16, -0.03)
Placebo (448)	2.16	-0.20	N.A.
Active Control [†] (Loratadine 10 mg) (180)	2.23	-0.45	-0.22 [‡] (-0.31, -0.13)

* Average of individual scores of nasal congestion, rhinorrhea, nasal itching, sneezing as assessed by patients on a 0-3 categorical scale.

[†] The study was not designed for statistical comparison between SINGULAIR and the active control (loratadine).

^{††} Statistically different from placebo (p=0.003); [‡] Statistically different from placebo (p≤0.001).

Figure 1



The study was not designed for statistical comparison between SINGULAIR and the active control (loratadine).

Paediatric patients 2 to 14 years of age

Efficacy studies have not been conducted in this age group.

5.2 PHARMACOKINETIC PROPERTIES

Absorption

Montelukast is rapidly and nearly completely absorbed following oral administration. For the 10 mg film-coated tablet, the mean peak plasma concentration (C_{max}) is achieved 3 hours (T_{max}) after administration in adults in the fasted state. The mean oral bioavailability is 64%. The oral bioavailability and C_{max} are not influenced by a standard meal.

For the 5 mg chewable tablet, the C_{max} is achieved in 2 hours after administration in adults in the fasted state. The mean oral bioavailability is 73% and is decreased to 63% by a standard meal. This is unlikely to have any clinical significance with chronic administration. Efficacy was demonstrated in clinical studies in children where the montelukast 5 mg chewable tablet was administered irrespective of food.

For the 4 mg chewable tablet, C_{max} is achieved 2 hours after the administration in paediatric patients 2 to 5 years of age in the fasted state.

The 4 mg oral granule formulation is bioequivalent to the 4 mg chewable tablet when administered to adults in the fasted state. The oral bioavailability is not influenced by a high-fat, high-calorie breakfast. This breakfast resulted in a longer t_{max} and a reduction in C_{max} of 36%. The 4 mg oral granule formulation is bioequivalent to the 4 mg chewable tablet when administered to fasted adults either alone or with applesauce.

The 10 mg film-coated tablets of montelukast are not bioequivalent to two 5 mg chewable tablets, and these two products should not be used interchangeably.

Distribution

Montelukast is more than 99% bound to plasma proteins. The steady-state volume of distribution of montelukast averages 8-11 liters. Studies in rats with radiolabeled montelukast indicate minimal distribution across the blood-brain barrier. In addition, concentrations of radiolabeled material at 24 hours postdose were minimal in all other tissues.

Metabolism

Montelukast is extensively metabolized. In studies with therapeutic doses, plasma concentrations of metabolites of montelukast are undetectable (<20 ng/mL) at steady state in adults and children. Cytochromes P450 2C8, and to a lesser extent 3A4 and 2C9, are involved in the metabolism of montelukast.

Excretion

The plasma clearance of montelukast averages 45 mL/min in healthy adults. Following an oral dose of radiolabeled montelukast, 86% of the radioactivity was recovered in 5-day fecal collections and <0.2% was recovered in urine. Coupled with estimates of montelukast oral bioavailability, this indicates that montelukast and its metabolites are excreted almost exclusively *via* the bile.

In several studies, the mean plasma half-life of montelukast ranged from 2.7 to 5.5 hours in healthy young adults. The pharmacokinetics of montelukast are nearly linear for oral doses up to 50 mg. During once-daily dosing with 10 mg montelukast, there is little accumulation of the parent drug in plasma (~14%).

5.3 PRECLINICAL SAFETY DATA

Genotoxicity

Montelukast sodium was found not to be genotoxic. Montelukast sodium was negative in microbial and mammalian cell mutagenesis assays, with and without metabolic activation.

There was no evidence of clastogenic activity in the *in vitro* chromosomal aberration assay in Chinese Hamster Ovary cells, with or without a microsomal enzyme activation system, or of DNA damage in the *in vitro* alkaline elution assay in rat hepatocytes. Similarly, there was no induction of chromosomal aberrations in bone marrow cells of male or female mice.

Carcinogenicity

Montelukast sodium was not carcinogenic when administered at oral doses of up to 200 mg/kg/day in 104 week study in rats, nor at oral doses up to 100 mg/kg/day in a 91 week study in mice. Systemic exposure in these studies, in terms of the plasma AUC for parent drug, was at least 30 times higher than that in humans at recommended dose levels.

6 PHARMACEUTICAL PARTICULARS

6.1 LIST OF EXCIPIENTS

Each 10 mg film-coated tablet contains the following inactive ingredients: microcrystalline cellulose, lactose monohydrate, croscarmellose sodium, hypromellose, and magnesium stearate. The film-coating consists of: hypromellose, hypromellose, titanium dioxide, iron oxide red, iron oxide yellow, and carnauba wax.

Each 4 mg and 5 mg chewable tablet contains the following inactive ingredients: mannitol, microcrystalline cellulose, hypromellose, croscarmellose sodium, artificial cherry flavour aromolok 181612 (Proprietary Ingredient ID 2916), magnesium stearate, aspartame, and iron oxide red.

Each packet of 4 mg oral granules contains the following inactive ingredients: mannitol, hypromellose, and magnesium stearate.

6.2 INCOMPATIBILITIES

Incompatibilities were either not assessed or not identified as part of the registration of these medicines.

6.3 SHELF LIFE

The expiry date can be found on the packaging. In Australia, information on the shelf life can be found on the public summary of the Australian Register of Therapeutic Goods (ARTG).

6.4 SPECIAL PRECAUTIONS FOR STORAGE

Store below 30°C. Protect from light and moisture.

6.5 NATURE AND CONTENTS OF CONTAINER

10 mg film-coated tablet - Supplied in blister packs of 4*, 14 and 28 tablets.

5 mg chewable tablet - Supplied in blister packs of 4* and 28 tablets.

4 mg chewable tablet - Supplied in blister packs of 4* and 28 tablets.

4 mg oral granules** - Supplied in packs of 28 packets.

* Supplied as starter packs only

** Not currently supplied in Australia

6.6 SPECIAL PRECAUTIONS FOR DISPOSAL

In Australia, any unused medicine or waste material should be disposed of by taking to your local pharmacy.

6.7 PHYSICOCHEMICAL PROPERTIES

SINGULAIR (montelukast sodium) is a selective and orally active leukotriene receptor antagonist that specifically inhibits the cysteinyl leukotriene CysLT₁ receptor.


SINGULAIR, (montelukast sodium) is described chemically as [R-(E)]-1-[[[1-[3-[2-(7-chloro-2-quinolinyl)ethenyl]phenyl]-3-[2-(1-hydroxy-1-methylethyl)phenyl]propyl]thio]methyl]cyclopropane acetic acid, monosodium salt.

The empirical formula is C₃₅H₃₅ClNaO₃S, and its molecular weight is 608.18.

Montelukast sodium is a hygroscopic, optically active, white to off-white, free-flowing powder. Montelukast sodium is freely soluble in ethanol, methanol, and water and practically insoluble in acetonitrile. Montelukast is the optically active R stereoisomer.

Chemical structure

The structural formula of montelukast sodium is:



CAS number

151767-02-1

7 MEDICINE SCHEDULE (POISONS STANDARD)

Prescription Only Medicine (Schedule 4)

8 SPONSOR

Merck Sharp & Dohme (Australia) Pty Limited
Level 1, Building A, 26 Talavera Road
Macquarie Park NSW 2113
www.msd-australia.com.au

9 DATE OF FIRST APPROVAL

11 May 1998 for 5 mg chewable tablets and 10 mg film-coated tablets

23 May 2001 for 4 mg chewable tablets

13 November 2002 for 4 mg oral granules

10 DATE OF REVISION

21 December 2018

Summary table of changes

Section Changed	Summary of new information
-----------------	----------------------------

All sections	PI reformat, minor editorial updates
4.7 Effects on Ability to drive and use machines	Revision of statement
4.8 Adverse Effects (Undesirable effects)	Addition of "obsessive compulsive symptoms" under post marketing subheading
6.1 List of Excipients	Revision to order of excipients