


KETOROLAC APOTEX Solution for Injection**NAME OF THE MEDICINE**

Ketorolac trometamol.

Chemical Name: 2-Amino-2-(Hydroxyethyl) propane-1, 3-diol (1*RS*)-5-benzoyl-2, 3-dihydro-1*H* pyrrolizine-1-carboxylate.

Structural Formula:



And enantiomer

WARNING

Ketorolac is a potent NSAID analgesic and the resulting NSAID-related adverse effects can be serious, for example gastrointestinal haemorrhage, surgical haemorrhage and renal impairment.

Increasing the dose of ketorolac beyond the recommendations in the product information will not provide better efficacy but will result in increasing risk of developing serious adverse effects.

Molecular Formula: $C_{19}H_{24}N_2O_6$

Molecular Weight: 376.41

CAS Registry Number: 74103-07-4

DESCRIPTION

Ketorolac Solution for Injection (ketorolac trometamol; synonym ketorolac tromethamine) is a non-narcotic analgesic belonging to the non-steroidal anti-inflammatory drug (NSAID) class of medicines with analgesic, anti-inflammatory and antipyretic properties.

Ketorolac Solution for Injection is a member of the pyrrolo-pyrrole group of NSAIDs. Ketorolac trometamol is 2-Amino-2-(Hydroxyethyl) propane-1, 3-diol (1*RS*)-5-benzoyl-2, 3-dihydro-1*H* pyrrolizine-1-carboxylate. The medicine is structurally and pharmacologically related to tolmetin and indomethacin, however, unlike these pyrrole acetic acid derivatives, ketorolac is a cyclic propionic derivative. Ketorolac solution for injection is a racemic mixture of [-] S and [+] R enantiomers, with the S-form having analgesic activity. Ketorolac is a white to off-white crystalline substance which discolours on exposure to light. It is soluble in water and has a pKa of 3.54.

Each ampoule contains ketorolac as the active ingredient as a sterile solution for intramuscular (IM) administration only. The injection solutions are clear and slightly yellow in colour. In addition, each ampoule contains the following inactive ingredients: ethanol, sodium chloride, water for injections and sodium hydroxide or hydrochloric acid to adjust pH.

PHARMACOLOGY

Pharmacological Actions

Ketorolac trometamol inhibits the cyclo-oxygenase enzyme system and hence synthesis of prostaglandins. It is considered to be a peripherally-acting analgesic. It does not have known effects on opiate receptors. It has no intrinsic effects on respiration and does not exacerbate opioid-related respiratory depression or sedation. Ketorolac trometamol has no significant CNS effects in animals and possesses no sedative or anxiolytic properties.

Pharmacokinetics

The pharmacokinetics of ketorolac in young healthy adult volunteers following single or multiple IM or recommended oral doses are linear.

Absorption

The IM form of ketorolac is rapidly and completely absorbed (100% bioavailable). Steady state plasma levels are achieved after dosing every 6 hours for one day. A mean peak plasma concentration of 2.2µg/mL occurs an average of 50 minutes after a single 30 mg dose.

Orally administered ketorolac is rapidly and completely absorbed with a peak plasma concentration of 0.52 to 1.31 µg/mL occurring 35 minutes after a single 10 mg dose in fasted subjects. A high fat diet decreased the rate, but not the extent of absorption, while antacid had no effect upon ketorolac absorption.

Distribution

More than 99% of ketorolac in plasma is protein bound. Even at high plasma concentrations (10 µg/mL) only approximately 5% of albumin binding sites will be occupied. Thus the unbound fraction for each enantiomer will be constant over the therapeutic range. A decrease in serum albumin, however, will result in increased free drug concentrations.

Plasma protein binding is independent of concentration. As ketorolac trometamol is a highly potent medicine and present in low concentrations in plasma, it would not be expected to displace other protein-bound medicines significantly.

Nearly all the medicine-related material circulating in plasma is ketorolac (96%) or the pharmacologically inactive p-hydroxyketorolac.

The mean apparent volume (V_{β}) of ketorolac trometamol following complete distribution is approximately 13 litres (this parameter was determined from single dose data).

Ketorolac trometamol poorly penetrates the blood brain barrier (levels in the cerebrospinal fluid were found to be 0.002 times or less than those in plasma).

Ketorolac crosses the placenta (mean umbilical/maternal vein concentration ratio for ketorolac was 0.116 and this ratio increased as the time from injection to sampling increased). Ketorolac has been detected in human milk at low concentrations (refer to PRECAUTIONS – Use in Pregnancy and Use in Lactation).

Metabolism

Ketorolac trometamol is largely metabolised in the liver. The major metabolic path of ketorolac in humans is glucuronic acid conjugation. P-hydroxylation is an additional minor pathway.

Excretion

The primary route of excretion of ketorolac and its metabolites (conjugates and a para-hydroxy metabolite) is in the urine (mean 91.4%) and the remainder (mean 6.1%) is excreted in the faeces.

The terminal plasma half-life of ketorolac is approximately in the range of 5-6 hours. No changes in clearance occur with chronic dosing.

Pharmacokinetics in Special Populations

Elderly Patients (65 years of age or older)

In the elderly, the mean terminal plasma half-life of ketorolac trometamol increases from 5 to 7 hours compared with young healthy volunteers (based on single dose data) (refer to DOSAGE AND ADMINISTRATION). There is little difference in the C_{max} for the two groups.

Renally-Impaired Patients

The mean half-life of ketorolac trometamol in renally-impaired patients is between 6 and 19 hours, and is dependent on the extent of the impairment (based on single dose data). There is poor correlation between creatinine clearance and total ketorolac trometamol clearance in the elderly and populations with renal impairment ($r = 0.5$). In patients with renal disease, the AUC increases by approximately 100% compared with healthy volunteers. The volume of distribution increases by up to 100%. The increase in volume of distribution of ketorolac trometamol implies an increase in unbound fraction. The AUC ratio of the ketorolac trometamol enantiomers in healthy subjects and patients remains similar, indicating there is no selective excretion of either enantiomer in patients compared to healthy subjects.

Hepatically-Impaired Patients

Patients with impaired hepatic function do not have any clinically important changes in ketorolac pharmacokinetics, although there is a statistically significant prolongation of T_{max} and terminal phase half-life compared to young healthy volunteers.

CLINICAL TRIALS

Intramuscular

Controlled clinical trials studying acute severe pain following major surgical procedures such as cholecystectomy, gastric bypass, abdominal hysterectomy, open reduction and fixation of fractures, lumbar laminectomy, and extraction of multiple impacted third molar teeth, have established the efficacy of ketorolac relative to other analgesics.

Given post-operatively, ketorolac 30 mg IM has an onset of action and peak analgesic efficacy comparable to morphine 12 mg IM or pethidine 100 mg IM, and is more effective than morphine 6 mg IM or pethidine 50 mg IM.

Ketorolac 30 mg IM has a longer duration of action than morphine 12 mg IM or pethidine 100 mg IM. Ketorolac 10 mg IM gives efficacy equal to or greater than morphine 6 mg IM or pethidine 50 mg IM.

Oral

Comparative controlled clinical trials studying acute pain following extraction of impacted third molars and orthopaedic, abdominal and gynaecological surgery have established the analgesic efficacy of oral ketorolac trometamol. In a series of clinical trials, TORADOL Tablets 10 mg have been found superior or equal to aspirin 650 mg, paracetamol 600 mg and 1000 mg, paracetamol 600 mg plus codeine 60 mg, paracetamol 1000 mg plus codeine 60 mg, and ibuprofen 400 mg.

INDICATIONS

Ketorolac is indicated for the short-term management of moderately severe, acute pain following surgical procedures. The total duration of ketorolac use should not exceed five days.

It is recommended that ketorolac parenteral be used in the immediate post-operative period. Patients can then be converted to the oral formulation (dependent on their analgesic needs), as outlined in the "DOSAGE AND ADMINISTRATION" Section (Refer to "Conversion from Parenteral to Oral Therapy"). The total period of treatment utilising the oral and/or intramuscular route of administration is not to exceed five days.

General

Ketorolac is not recommended for use as an obstetrical pre-operative medication or for obstetrical analgesia because it has not been adequately studied for use in these circumstances and because the known effects of drugs that inhibit prostaglandin biosynthesis on uterine contraction and foetal circulation.

There is no satisfactory evidence for the use of ketorolac in acute exacerbations of chronic painful inflammatory conditions (e.g. rheumatoid or osteoarthritis).

CONTRAINDICATIONS

Ketorolac is contraindicated in patients:

- With severe heart failure (refer to PRECAUTIONS – Heart Failure).
- With dehydration or hypovolaemia from any other cause.
- With moderate or severe renal impairment (serum creatinine > 180 micromol/L), or in patients at risk of renal failure due to volume depletion or dehydration (refer to PRECAUTIONS – Renal Effects).
- With active, or a history of gastrointestinal bleeding or perforation, related to previous NSAIDs therapy. Active, or history of recurrent peptic ulcer/haemorrhage (two or more distinct episodes of proven ulceration or bleeding) (refer to PRECAUTIONS – Gastrointestinal Effects).
- With a history of haemorrhagic diatheses, including coagulation disorders (refer to PRECAUTIONS – Haematologic Effects).
- Who have surgery with a high risk of haemorrhage or incomplete haemostasis; and those at high risk of bleeding (refer to PRECAUTIONS – Haematologic Effects).
- With suspected or confirmed cerebrovascular (intracranial) bleeding (refer to PRECAUTIONS – Haematologic Effects).
- On anticoagulation therapy (refer to PRECAUTIONS – Interactions with Other Medicines).
- Receiving aspirin, other NSAIDs, oxpentifylline, probenecid, or lithium (refer to PRECAUTIONS – Interactions with other Medicines).
- With hypersensitivity to ketorolac trometamol or other NSAIDs and those patients in whom aspirin or other prostaglandin synthetase inhibitors induce allergic reactions. Severe anaphylactic-like reactions have been observed in such patients. If such symptoms occur during therapy, treatment should be discontinued (refer to PRECAUTIONS – Anaphylactic Reactions).
- With the complete or partial syndrome of nasal polyps, angioedema or bronchospasm (refer to PRECAUTIONS – Anaphylactic Reactions).
- with a history of asthma (refer to PRECAUTIONS – Anaphylactic Reactions).
- With prior history of Stevens-Johnson syndrome or vesicular bullous rash (refer to PRECAUTIONS – Severe Skin Effects and ADVERSE EFFECTS).

Ketorolac is also contraindicated for:

- Neuraxial (epidural or intrathecal) administration due to the alcohol content of the solution for injection.
- Prophylactic administration before surgery, due to inhibition of platelet aggregation; and intraoperatively because of the increased risk of bleeding.
- Use in pregnancy, labour delivery or lactation (refer to PRECAUTIONS – Use in Pregnancy and Use in Lactation).
- Children under 16 years-of-age.

PRECAUTIONS

Effects on Fertility

Ketorolac may impair fertility and is not recommended in women attempting to conceive.

Use in Pregnancy (Category C)

Ketorolac is not recommended for use during pregnancy, labour and delivery (refer to CONTRAINDICATIONS).

NSAIDs inhibit prostaglandin synthesis and, when given during the latter part of pregnancy, may cause closure of the foetal ductus arteriosus, foetal renal impairment, inhibition of platelet aggregation, and delay labour and birth. Continuous treatment with NSAIDs during the last trimester of pregnancy should only be given on sound indications. During the last few days before expected birth, agents with an inhibitory effect on prostaglandin synthesis should be avoided.

Reproductive studies with ketorolac have been performed in rabbits and rats at oral doses of 3.6 and 10 mg/kg/day, respectively. The results from these studies did not reveal any significant evidence of harm to the foetus. However, studies in rabbits have shown a small increase in the incidence of major vessel anomalies.

Use in Lactation

Ketorolac is not recommended for treatment of nursing mothers (refer to CONTRAINDICATIONS).

After a single oral administration of 10 mg of ketorolac to humans, the maximum milk concentration observed was 7.3 ng/mL and the maximum milk-to-plasma ratio was 0.037. After one day of dosing (qid), the maximum milk concentration was 7.9 ng/mL and the maximum milk-to-plasma ratio was 0.025.

Paediatric Use

The safety and efficacy in children have not been established, therefore, ketorolac is not recommended for use in children under 16 years of age (refer to CONTRAINDICATIONS).

Use in the Elderly

Because ketorolac trometamol is cleared somewhat more slowly by the elderly (refer to PHARMACOKINETICS) who are also more sensitive to the fluid retaining, gastrointestinal toxicity and renal impairment effects (refer to PRECAUTIONS) of this medicine, extra caution and the lowest effective dose should be used when treating the elderly with ketorolac (refer to DOSAGE AND ADMINISTRATION).

Ability to Drive and Operate Machinery

Some patients may experience drowsiness, dizziness, vertigo, insomnia or depression with the use of ketorolac. If patients experience these, or other similar undesirable effects, they should exercise caution in carrying out activities that require alertness.

Cardiovascular Thrombotic Events

Observational studies have indicated that non-selective NSAIDs may be associated with an increased risk of serious cardiovascular events, including myocardial infarction and stroke, which may increase with dose or duration of use. Patients with cardiovascular disease or cardiovascular risk factors may also be at greater risk. To minimise the potential risk of an adverse cardiovascular event in patients taking an NSAID, especially in those with cardiovascular risk factors, the lowest effective dose should be used for the shortest possible duration (refer to DOSAGE AND ADMINISTRATION). Although ketorolac has not shown to increase thrombotic events such as myocardial infarction, there are insufficient data to exclude such a risk for ketorolac.

There is no consistent evidence that the concurrent use of aspirin mitigates the possible increased risk of serious cardiovascular thrombotic events associated with NSAID use.

Hypertension

NSAIDs may lead to the onset of new hypertension or worsening of pre-existing hypertension and patients taking anti-hypertensives with NSAIDs may have an impaired anti-hypertensive response. Caution is advised when prescribing NSAIDs to patients with hypertension. Blood pressure should be monitored closely during initiation of NSAID treatment and at regular intervals thereafter.

Heart Failure

Fluid retention and oedema have been observed in some patients taking NSAIDs; therefore caution is advised in patients with fluid retention, cardiac decompensation, heart failure, hypertension or similar conditions.

Gastrointestinal Effects

Ketorolac can cause gastrointestinal irritation, ulcers, perforation or bleeding, which can be fatal, at any time, with or without warning symptoms or a previous history of serious gastrointestinal events.

Upper gastrointestinal ulcers, gross bleeding or perforation caused by NSAIDs occur in approximately 1% of patients treated for 3 - 6 months and in about 2 - 4% of patients treated for one year. These trends continue with longer duration of use, increasing the likelihood of developing a serious gastrointestinal effect at some time during the course of therapy. However, even short-term therapy is not without risk.

The risk of gastrointestinal bleeding, ulceration or perforation increases with dose and duration of treatment; in patients with a history of ulcer, particularly if complicated with haemorrhage or perforation; in the elderly; and in those with a history of smoking or alcoholism. Caution is advised in these patients and treatment should commence on the lowest dose available. Combination therapy with gastro-protective agents (e.g. misoprostol or proton-pump inhibitors) should be considered for these patients.

Elderly and debilitated individuals are more susceptible to gastrointestinal complications (refer to DOSAGE AND ADMINISTRATION - Elderly for dosage reductions in this patient group). Most reports of fatal gastrointestinal effects are in this population.

NSAIDs should be given with care to patients with a history of inflammatory bowel disease (ulcerative colitis; Crohn's disease) as their condition may be exacerbated.

Caution should be advised in patients receiving concomitant medications which could increase the risk of gastrointestinal ulceration or bleeding, such as NSAIDs, oral corticosteroids; anticoagulants such as warfarin; selective serotonin reuptake inhibitors (SSRIs); or anti-platelet agents such as aspirin, as these combinations may increase the risk of serious gastrointestinal adverse effects (refer to PRECAUTIONS - Interactions with Other Medicines). Combination therapy with gastro-protective agents (e.g. misoprostol or proton-pump inhibitors) should be considered for these patients.

Prescribers should warn patients about the signs and symptoms of serious gastrointestinal toxicity. Patients administered ketorolac should be instructed to advise their physician immediately if they experience any unusual abdominal symptoms (especially gastrointestinal bleeding). Ketorolac should be discontinued, appropriate treatment instituted and the patient closely monitored.

In a non-randomised, in-hospital post-marketing surveillance study, increased rates of clinically serious gastrointestinal bleeding were seen in patients 65 years-of-age and under who received an average daily dose of greater than 90 mg of ketorolac administered intramuscularly as compared to those patients receiving parenteral opioids.

Renal Effects

As with other NSAIDs that inhibit prostaglandin biosynthesis, elevations of serum urea, nitrogen, potassium and creatinine have been reported in clinical trials with ketorolac, and can occur after one dose. Ketorolac trometamol and its metabolites are eliminated primarily by the kidneys which, in patients with reduced creatinine clearance, will result in diminished clearance of the medicine.

Patients with moderate to severe impairment of renal function should not receive ketorolac (refer to DOSAGE AND ADMINISTRATION - Mild Renal Impairment for dosage reduction in patients with mild renal impairment i.e. serum creatinine < 180 micromol/L). Ketorolac should be used with caution in patients with a history of kidney disease. Renal function should be monitored in patients who have had more than a single intramuscular dose of ketorolac, particularly in elderly patients.

As with other NSAIDs that inhibit prostaglandin biosynthesis, the following renal abnormalities may be associated with the use of ketorolac: glomerular nephritis, interstitial nephritis, renal papillary necrosis, nephrotic syndrome, acute renal failure and hyperkalaemia. Other renal conditions/diseases are possible.

Caution should be observed in patients with conditions leading to a reduction in blood volume and/or renal blood flow, where renal prostaglandins have a supportive role in the maintenance of renal perfusion. In these patients, administration of an NSAID may cause a dose-dependent reduction in renal prostaglandin formation and may precipitate overt renal failure. Patients at greatest risk of this effect are those who are volume depleted because of blood loss or severe dehydration,

patients with impaired renal function, heart failure, liver dysfunction, the elderly and those taking diuretics. Discontinuation of NSAID therapy is typically followed by recovery to the pre-treatment state. Inadequate fluid/blood replacement during surgery, leading to hypovolaemia (refer to CONTRAINDICATIONS), may lead to renal dysfunction which could be exacerbated when ketorolac is administered. Therefore, volume depletion should be corrected and close monitoring of serum urea, serum creatinine and urine output is recommended until the patient is normovolaemic. In patients on renal dialysis, ketorolac clearance is reduced to approximately half the normal rate and terminal half-life increases approximately three-fold.

Haematologic Effects

Ketorolac inhibits platelet aggregation and may prolong bleeding time. Unlike the prolonged effects from aspirin, the inhibition of platelet function by ketorolac resolves within 24 to 48 hours after the medicine is discontinued. Ketorolac does not affect platelet count, prothrombin time (PT) or partial thromboplastin time (PTT). In controlled clinical studies, the incidence of clinically significant post-operative bleeding was 5/1170 (0.4%) compared to 1/570 (0.2%) in the control groups receiving opiates.

The use of ketorolac in patients who have coagulation disorders should be undertaken very cautiously, and those patients carefully monitored. PATIENTS ON ANTICOAGULATION THERAPY (e.g., HEPARIN OR WARFARIN) MAY BE AT INCREASED RISK OF BLEEDING IF GIVEN KETOROLAC CONCURRENTLY (refer to CONTRAINDICATIONS). The concomitant use of ketorolac and prophylactic low-dose heparin has not been studied extensively and may also be associated with an increased risk of bleeding. Concomitant administration of dextrans may also increase the risk of post-operative bleeding. Patients receiving other medicines that affect haemostasis should be carefully observed if ketorolac is administered (refer to PRECAUTIONS – Interactions with Other Medicines).

In post-marketing experience, post-operative wound haemorrhage has been reported in association with the immediate peri-operative use of intramuscular ketorolac. Therefore, ketorolac should not be used in patients who have had surgery with a high risk of haemorrhage or incomplete haemostasis. Caution should be used where strict haemostasis is critical, e.g. in cosmetic or day care surgery, resection of the prostate or tonsillectomy. Haematomata, other signs of wound haemorrhage and epistaxis have been reported with the use of ketorolac. Physicians should be aware of the pharmacological similarity of ketorolac to other NSAIDs that inhibit cyclo-oxygenase and the risk of bleeding, particularly in the elderly.

Hepatic Effects

As with other NSAIDs, borderline elevations of one or more liver function tests may occur in up to 15% of patients. These abnormalities may progress, may remain essentially unchanged, or may be transient with continued therapy. The ALT (SGPT) test is probably the most sensitive indicator of liver injury. Meaningful (3 times the upper limit of normal) elevations of ALT or AST (SGOT) have been reported in controlled clinical trials (with the oral formulation of ketorolac trometamol) in less than 1% of patients. If clinical signs and symptoms consistent with liver disease develop, or if systemic manifestations occur (e.g., eosinophilia, rash etc.), ketorolac should be discontinued.

Patients with impaired hepatic function from cirrhosis do not have any clinically important changes in ketorolac clearance. Studies to assess the pharmacokinetics of ketorolac in patients with active hepatitis or cholestasis have not been done. However, a patient with symptoms and/or signs suggesting liver dysfunction, or in whom an abnormal liver test has occurred, should be evaluated for evidence of the development of a more severe hepatic effects while on therapy with ketorolac.

Severe Skin Effects

NSAIDs may very rarely cause serious cutaneous adverse effects such as exfoliative dermatitis, Stevens-Johnson Syndrome (SJS) and toxic epidermal necrolysis (TEN), which can be fatal and occur without warning. These serious adverse effects are idiosyncratic and are independent of dose or duration of use. Patients should be advised of the signs and symptoms of serious skin effects and to consult their physician at the first appearance of a skin rash or other sign of hypersensitivity.

Injection Site Effects

Ketorolac Injection administered intramuscularly has produced pain at the injection site in 2 to 4% of patients. Ecchymosis, bruising, haematoma and tingling at the injection site have rarely been reported. Adverse local effects may be minimised by applying pressure at the injection site for 15 to 30 seconds after administration. There has been no evidence (e.g. alterations in serum creatine kinase [CK] or

creatine phosphokinase [CPK] concentrations) of substantial adverse muscular tissue effects following single or multiple intramuscular injections of ketorolac.

Anaphylactic Reactions

Anaphylactic (anaphylactoid) reactions (including, but not limited to, anaphylaxis, bronchospasm, flushing, rash, hypotension, laryngeal oedema and angioedema) may occur in individuals with or without a history of hypersensitivity to aspirin, other NSAIDs, or ketorolac. These may also occur in individuals with a history of angioedema, bronchospastic reactivity (e.g. asthma) and nasal polyps. Anaphylactoid reactions, like anaphylaxis, may have a fatal outcome. Therefore, ketorolac should be used with caution in patients with a history of asthma and in patients with the complete or partial syndrome of nasal polyps, angioedema and bronchospasm.

Special Senses

Adverse ophthalmological effects have been observed with non-steroidal anti-inflammatory agents; accordingly, patients who develop visual disturbances during treatment with ketorolac should have an ophthalmological examination.

Drug Abuse and Physical Dependence

Ketorolac is not a narcotic agonist or antagonist. Subjects did not show any subjective symptoms or objective signs of drug withdrawal upon abrupt discontinuation of intramuscular dosing. Ketorolac did not exhibit activity in classical animal studies which are reasonable predictors of opiate analgesic action (hot plate and tail withdrawal test). *In vitro* ketorolac does not bind to opiate receptors. These studies demonstrate that ketorolac does not have central opiate-like activity.

General

Undesirable effects may be minimised by using the lowest minimum effective dose for the shortest duration necessary to control symptoms (refer to DOSAGE AND ADMINISTRATION and PRECAUTIONS).

Ketorolac is not an anaesthetic agent and possesses no sedative or anxiolytic properties, therefore it is not recommended as a pre-operative or intra-operative medication for the support of anaesthesia when these effects are required. Ketorolac IM should not be used for spinal or epidural administration. Ketorolac IM contains ethanol 10% (refer to CONTRAINDICATIONS).

The total duration of ketorolac treatment should not exceed five days.

INTERACTIONS WITH OTHER MEDICINES

Protein Binding

Ketorolac is highly bound to human plasma protein (mean 99.2%) and binding is independent of concentration. The *in vitro* binding of warfarin to plasma proteins is only slightly reduced by ketorolac (99.5% control vs. 99.3% binding with ketorolac concentrations of 5 to 10 µg/mL). Ketorolac does not alter digoxin protein binding.

In vitro studies indicated that at therapeutic concentrations of salicylate (300 µg/mL), the binding of ketorolac was reduced from approximately 99.2% to 97.5%. Therapeutic concentrations of digoxin, warfarin, ibuprofen, naproxen, paracetamol, phenytoin, tolbutamide and piroxicam did not alter ketorolac protein binding. Because ketorolac is a highly potent medicine and present in low concentrations in plasma, it would not be expected to displace other protein bound medicines significantly.

Enzyme Induction / Inhibition

There is no evidence in animal or human studies that ketorolac induces or inhibits the hepatic enzymes capable of metabolising itself or other medicines. Hence, ketorolac would not be expected to alter the pharmacokinetics of other medicines due to enzyme induction or inhibition mechanisms.

Warfarin

The concurrent use of NSAIDs and warfarin has been associated with severe, sometimes fatal, haemorrhage. The exact mechanism of the interaction between NSAIDs and warfarin is unknown, but may involve enhanced bleeding from NSAID-induced gastrointestinal ulceration, or an additive effect

of anticoagulation by warfarin and inhibition of platelet function by NSAIDs. Ketorolac should be used in combination with warfarin only if absolutely necessary, and patients taking this combination of medicines should be closely monitored.

Anticoagulant Therapy and Other Drugs Affecting Haemostasis

NSAIDs may enhance the effects of anti-coagulants, such as warfarin, low-molecular weight heparin and dextrans. (refer to CONTRAINDICATIONS and PRECAUTIONS - Haematologic Effects). Unlike the prolonged effects from aspirin, platelet function returns to normal within 24 – 48 hours after ketorolac is discontinued.

Aspirin and Other NSAIDs

In patients concurrently receiving aspirin or other NSAIDs, the risk of inducing serious NSAID-related adverse effects may be increased (refer to CONTRAINDICATIONS).

Oxpentifylline

When ketorolac is administered concurrently with oxpentifylline, there is an increased tendency to bleeding (refer to CONTRAINDICATIONS).

Probenecid

Concomitant administration of oral ketorolac and probenecid resulted in decreased clearance of ketorolac and significant increases in ketorolac plasma levels (total AUC increased approximately 3-fold from 5.4 to 17.8 µg/h/mL) and terminal half-life increased approximately 2-fold from 6.6 to 15.1 hours. Therefore the concomitant use of ketorolac and probenecid is contraindicated.

Lithium

Inhibition of renal lithium clearance, leading to an increase in plasma lithium concentration, has been reported with some prostaglandin synthesis-inhibiting medicines. The effect of ketorolac on plasma lithium has not been studied, but cases of increased plasma levels during ketorolac therapy have been reported.

Methotrexate

Concomitant administration of methotrexate and some NSAIDs has been reported to reduce the clearance of methotrexate, enhancing the toxicity of methotrexate. The effect of ketorolac on methotrexate clearance has not been studied.

ACE-Inhibitors

As with other NSAIDs, ketorolac may increase the risk of renal impairment associated with the use of ACE-inhibitors, particularly in patients that are actually or effectively volume depleted.

Combination use of ACE inhibitors or angiotensin receptor antagonists, anti-inflammatory medicines and thiazide diuretics (Triple Whammy)

The use of an ACE inhibiting medicine (ACE-inhibitor or angiotensin receptor antagonist), an anti-inflammatory medicine (NSAID or COX-2 inhibitor) and a thiazide diuretic at the same time (Triple Whammy) increases the risk of renal impairment. This includes use in fixed-combination products containing more than one class of medicine. Combined use of these medications should be accompanied by increased monitoring of serum creatinine, particularly during initiation of the combination. The combination of medicines from these three classes should be used with caution particularly in elderly patients or those with pre-existing renal impairment.

Diuretics

Ketorolac reduces the diuretic response to frusemide in normovolaemic healthy subjects by approximately 20%.

Nephrotoxic Agents

The use of medicines with nephrotoxic activity (e.g. aminoglycoside antibiotics) should be avoided when using ketorolac.

There is an increased risk of gastrointestinal bleeding when anti-platelet agents and SSRIs are combined with NSAIDs.

Selective Serotonin Reuptake Inhibitors (SSRIs)

There is an increased risk of gastrointestinal bleeding when anti-platelet agents and SSRIs are combined with NSAIDs.

Antiepileptic Medicines

Sporadic cases of seizures have been reported during concomitant use of ketorolac and antiepileptic medicines (phenytoin, carbamazepine).

Psychoactive Medicines

Hallucinations have been reported when ketorolac was used in patients taking psychoactive medicines (fluoxetine, thiothixene, alprazolam).

ADVERSE EFFECTS

Parenteral Administration

Physicians using ketorolac Injection should be alert for the usual complications of non-steroidal anti-inflammatory treatment.

The adverse effects listed below were reported to be probably related to ketorolac in clinical trials in which patients received up to 20 doses, in five days, of **post-operatively** administered ketorolac 30 mg intramuscularly and in clinical trials in which patients received up to 8 doses in two days, of **post-operatively** administered ketorolac 30 mg intravenously.

Incidence between 3% and 9%:

Gastrointestinal Disorders: nausea, dyspepsia, gastrointestinal pain.

Nervous System Disorders: drowsiness.

Incidence between 1% and 3%:

Gastrointestinal Disorders: diarrhoea.

General Disorders and Administration Site Conditions: oedema, injection site pain was reported by 2% of patients in multi-dose studies (compared with 5% for the morphine control group).

Nervous System Disorders: dizziness, headache.

Skin and Subcutaneous Tissue Disorders: sweating.

Incidence 1% or less:

Ear Disorders: vertigo.

Eye disorders: abnormal vision.

Gastrointestinal Disorders: constipation, flatulence, gastrointestinal fullness, liver function abnormalities, melaena, peptic ulcer, rectal bleeding, stomatitis, vomiting.

General Disorders and Administration Site Conditions: asthenia, excessive thirst.

Musculoskeletal and Connective Tissue Disorders: myalgia.

Nervous System Disorders: dry mouth, paraesthesia, stimulation, abnormal taste.

Psychiatric Disorders: nervousness, abnormal thinking, depression, euphoria, insomnia, inability to concentrate.

Renal and Urinary Disorders: increased urinary frequency, oliguria.

Respiratory, Thoracic and Mediastinal Disorders: dyspnoea, asthma.

Skin and Subcutaneous Tissue Disorders: pruritus, urticaria, purpura.

Vascular Disorders: vasodilation, pallor.

Post-marketing Adverse Effects

The following international post-marketing adverse effects, although rare, have been reported spontaneously for patients who have received ketorolac trometamol:

Blood and Lymphatic System Disorders: thrombocytopenia, epistaxis, haematoma, angioedema.

Cardiac Disorders: bradycardia, palpitations, cardiac failure.

Ear Disorders: hearing loss, tinnitus, vertigo.

Eye Disorders: abnormal vision.

Gastrointestinal Disorders: gastrointestinal haemorrhage, peptic ulceration, gastrointestinal perforation, nausea, vomiting, diarrhoea, flatulence, eructation, constipation, dyspepsia, abdominal pain/discomfort, melaena, haematemesis, stomatitis, ulcerative stomatitis, oesophagitis, rectal bleeding, pancreatitis, dry mouth, fullness, exacerbation of colitis and Crohn's disease, gastritis.

General Disorders and Administration Site Conditions: weight gain, injection site reactions, pallor, fever, asthenia, oedema, excessive thirst, chest pain.

Hepatobiliary Disorders: hepatitis, cholestatic jaundice, liver failure.

Immune System Disorders: anaphylaxis, anaphylactoid reactions, hypersensitivity reactions such as flushing, rash, hypotension, bronchospasm and laryngeal oedema (refer to PRECAUTIONS – Anaphylactic Reactions).

Infection: infection, aseptic meningitis.

Investigations: abnormal liver function tests, increased serum urea, increased creatinine, prolonged bleeding time.

Metabolic and Nutrition Disorders: hyponatraemia, hyperkalaemia, anorexia.

Musculoskeletal and Connective Tissue Disorders: myalgia.

Nervous System Disorders: headache, dizziness, paraesthesia, convulsions, extrapyramidal symptoms, hyperkinesia, taste abnormality, dry mouth, drowsiness.

Psychiatric Disorders: abnormal dreams, hallucinations, anxiety, psychotic reactions, abnormal thinking, depression, insomnia, nervousness, euphoria, impaired concentration ability.

Renal and Urinary Disorders: acute renal failure, urinary retention, increased urinary frequency, interstitial nephritis, nephrotic syndrome, haemolytic uraemic syndrome, oliguria, flank pain with or without haematuria and/or azotemia (refer to PRECAUTIONS – Renal Effects).

Reproductive System and Breast Disorders: female infertility.

Respiratory, Thoracic and Mediastinal Disorders: asthma, dyspnoea, pulmonary oedema.

Skin and Subcutaneous Tissue Disorders: rash, Stevens-Johnson syndrome (SJS), exfoliative dermatitis, toxic epidermal necrolysis (TEN), maculopapular rash, Lyell's syndrome, pruritus, urticaria, purpura, sweating.

Vascular Disorders: hypotension, hypertension, flushing, pallor, post-operative wound haemorrhage (rarely requiring blood transfusion).

DOSAGE AND ADMINISTRATION

Ketorolac dosage should be adjusted according to the severity of the pain and the response of the patient. **The lowest effective dose should be used for the shortest possible time in all patient populations.**

Opiate analgesics (e.g. morphine and pethidine) may be used concomitantly, and may be required for optimal analgesic effect in the early post-operative period when pain is most severe, or when the anxiolytic effects and/or sedative effects of opiates are desired. Ketorolac has been administered with morphine in several clinical trials of post-operative pain without evidence of adverse interactions. Ketorolac does not exacerbate opioid-related respiratory depression or sedation. When used in association with intramuscular ketorolac the daily dose of opioid is usually less than that which is normally required.

Hypovolaemia should be corrected prior to administration of ketorolac.

The total duration of ketorolac administration should not exceed 5 days because adverse effects may increase with prolonged usage.

Intramuscular Administration

The intramuscular injection should be given slowly and deeply into the muscle.

Adults (under 65 years of age)

The usual recommended initial intramuscular dose is 10 mg to 30 mg, followed by 10 mg to 30 mg at 4 to 6 hourly intervals, up to a maximum daily dose of 90 mg.

Elderly (65 years of age and older)

An initial intramuscular dose of 10 mg to 15 mg, followed by 10 mg to 15 mg at 4 to 6 hourly intervals, up to a maximum daily dose of 60 mg.

Mild Renal Impairment

If ketorolac is used in patients with mildly impaired renal function (serum creatinine values: males between 130 and 180 micromol/L; females between 120 and 180 micromol/L) the lower end of the intramuscular dosage range should be used. The total daily dose should not exceed 60 mg. Ketorolac is contraindicated in patients with more severe degrees of renal impairment (refer to CONTRAINDICATIONS).

Other

For patients under 50 kg in body weight or for patients with less severe pain, the lower end of the intramuscular ketorolac dosage range is recommended. The total daily dose should not exceed 60 mg.

Ketorolac solution for injection is for single use in one patient only. Discard any residue.

Conversion from Intramuscular to Oral Therapy

For patients being converted from intramuscular ketorolac to oral ketorolac, the total combined daily dose should not exceed 90 mg (60 mg for the elderly, mild renally-impaired patients and patients weighing less than 50 kg) and the oral component should not exceed 40 mg (30 mg to 40 mg for the elderly) on the day the change of formulation is made. Patients should be converted to oral treatment as soon as possible.

This brand of ketorolac does not include an oral presentation, therefore patients who need to be converted to oral therapy should be converted onto another brand.

OVERDOSAGE

Symptoms

Single overdoses of ketorolac have been variously associated with abdominal pain, nausea, vomiting, hyperventilation, peptic ulcers and/or erosive gastritis and renal dysfunction which have resolved after discontinuation of dosing.

Gastrointestinal bleeding may occur. Hypertension, acute renal failure, respiratory depression and coma may occur after the ingestion of NSAIDs but are rare.

Anaphylactoid reactions have been reported with therapeutic ingestion of NSAIDs and may occur following an overdose.

Treatment

Patients should be managed by symptomatic and supportive care following NSAID overdose. There are no specific antidotes. Dialysis does not significantly clear ketorolac from the blood stream.

For information on the management of overdose, contact the Poison Information Centre on 131126 (Australia).

PRESENTATION AND STORAGE CONDITIONS

KETOROLAC APOTEX Solution for Injection is intended for intramuscular administration.

Each solution for injection ampoule contains 10mg or 30mg of ketorolac as the active ingredient.

10 mg/mL Solution for Injection:

Clear and slightly yellow in colour solution.

1 mL solution in clear glass ampoules, in packs of 5 (AUST R 217074).

30 mg/mL Solution for Injection:

Clear and slightly yellow in colour solution.

1 mL solution in clear glass ampoules, in packs of 5 (AUST R 217079).

Not all strengths, pack types and/or pack sizes may be available.

Storage

Store below 30 °C. Protect from light.

NAME AND ADDRESS OF THE SPONSOR

Apotex Pty Ltd
16 Giffnock Avenue
Macquarie Park NSW 2113

APO and APOTEX are registered trademarks of Apotex Inc.

POISON SCHEDULE OF THE MEDICINE

S4 – Prescription Only Medicine.

Date of first inclusion in the Australian Register of Therapeutic Goods (the ARTG): 27-Oct-2014