

# TOPROL-XL™

**23.75, 47.5, 95, 190 TABLETS**

Metoprolol succinate

---

## **Consumer Medicine Information**

### **What is in this leaflet**

This leaflet answers some of the common questions people ask about TOPROL-XL tablets. It does not contain all the information that is known about TOPROL-XL.

It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor will have weighed the risks of you taking TOPROL-XL against the benefits they expect it will have for you.

**If you have any concerns about taking this medicine, ask your doctor or pharmacist.**

**Keep this leaflet with the medicine.**

You may need to read it again.

## **What TOPROL-XL is used for**

TOPROL-XL belongs to a group of medicines called beta-blockers.

TOPROL-XL tablets are used to treat heart failure. It is used in combination with other medicines to treat your condition.

It works by affecting the body's response to some nerve impulses, especially in the heart. As a result, it decreases the heart's need for blood and oxygen and therefore reduces the amount of work the heart has to do. It also helps the heart to beat more regularly.

**Ask your doctor if you have any questions about why TOPROL-XL has been prescribed for you.**

Your doctor will have explained why you are being treated with TOPROL-XL and told you what dose to take.

Your doctor may have prescribed this medicine for another reason.

**Follow all directions given to you by your doctor carefully.**

They may differ from the information contained in this leaflet.

TOPROL-XL is not addictive.

# **Before you take TOPROL-XL**

## **When you must not take it**

**Do not take TOPROL-XL tablets if:**

- **you have any allergies to metoprolol succinate, the active ingredient in TOPROL-XL tablets, or any of the ingredients listed at the end of this leaflet, or any other beta-blocker medicine.**

Some of the symptoms of an allergic reaction may include shortness of breath, wheezing or difficulty breathing; swelling of the face, lips, tongue or other parts of the body; rash, itching or hives on the skin or you may feel faint.

- **you have or have had asthma difficulty in breathing, wheezing, bronchitis or other lung problems in the past**
- **you have a history of allergic problems, including hayfever**
- **you have low blood pressure**
- **you have a very slow heartbeat (less than 45-50 beats/minute)**
- **you have certain other heart conditions**
- **you have phaeochromocytoma (a rare tumour of the adrenal gland) which is not being treated already with other medicines**

- **you have a severe blood vessel disorder causing poor circulation in the arms and legs**
- **you are receiving/having emergency treatment for shock or severely low blood pressure.**

**If you are not sure whether any of these apply to you, check with your doctor.**

**Do not use TOPROL-XL if the expiry date (EXP) printed on the pack has passed or if the packaging is torn or shows signs of tampering.**

If it has expired or is damaged, return it to your pharmacist for disposal.

**Do not give TOPROL-XL to children.**

The safety and effectiveness of TOPROL-XL in children has not been established.

**If you are not sure whether you should start taking this medicine, talk to your doctor.**

**Before you start to take it**

**Tell your doctor if you have any allergies to:**

- metoprolol succinate or any of the ingredients listed at the end of this leaflet
- any other medicine including other beta-blocker medicines

- any other substances, such as foods, preservatives or dyes

**Tell your doctor if you have, or have had, any medical conditions, especially the following:**

- asthma or other lung problems, even if you have had them in the past
- allergic problems, including hayfever
- diabetes
- very slow heart beat (less than 45-50 beats/minute)
- severe blood vessel disorder causing poor circulation in the arms and legs
- liver problems
- kidney problems
- certain types of angina
- any other heart problems
- phaeochromocytoma, a rare tumour of the adrenal gland
- hyperthyroidism (an overactive thyroid gland)

**Tell your doctor, if you are pregnant or intend to become pregnant.**

Like most beta- blocker medicines, TOPROL-XL is not recommended for use during pregnancy.

## **Tell your doctor, if you are breast-feeding or plan to breast-feed**

The active ingredient in TOPROL-XL passes into breast milk and therefore there is a possibility that the breast-fed baby may be affected.

**If you have not told your doctor about any of these things, tell them before you take TOPROL-XL.**

## **Taking other medicines**

**Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.**

Some medicines and TOPROL-XL may interfere with each other. These include:

- other beta-blocker medicines, including beta-blocker eye drops
- calcium channel blockers or calcium antagonists, medicines used to treat high blood pressure and angina, for example verapamil and diltiazem
- medicines used to treat high blood pressure, for example clonidine, hydralazine, and prazosin
- medicines used to treat abnormal or irregular heartbeat, for example amiodarone, disopyramide and quinidine

- medicines used to treat arthritis, pain, or inflammation, for example indomethacin and ibuprofen
- digoxin, a medicine used to treat heart failure
- medicines used to treat diabetes
- medicines used to treat bacterial infections, for example rifampicin
- cimetidine, a medicine used to treat stomach ulcers
- medicines used to treat depression
- warfarin, a medicine used to prevent blood clots
- monoamine-oxidase inhibitors (MAOIs)

These medicines may be affected by TOPROL-XL or may affect how well it works. You may need to take different amounts of your medicine, or you may need to take different medicines. Your doctor will advise you.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking TOPROL-XL.

**If you have not told your doctor about any of these things, tell them before you take any TOPROL-XL.**

## **How to take TOPROL-XL**

**Follow all directions given to you by your doctor or pharmacist carefully.**

They may differ from the information contained in this leaflet.

**If you do not understand the instructions on the pack, ask your doctor or pharmacist for help.**

## **How much to take**

The usual starting dose is half a 23.75 mg or a whole 23.75 mg tablet once a day for one to two weeks. The dose is then usually doubled every second week up to a maximum dose of 190 mg once daily or to the highest tolerated dose.

**TOPROL-XL tablets may be broken in half or swallowed whole with a glass of water as directed by your doctor. TOPROL-XL tablets should not be chewed or crushed.**

If you are taking other medicines your doctor may need to change the dose of them to obtain the best results for you.

## **When to take TOPROL-XL**

Take your medicine at about the same time each day, preferably together with the morning meal.

Taking it at the same time each day will have the best effect. It will also help you remember when to take it.

## **How long to take TOPROL-XL**

Continue taking your medicine for as long as your doctor tells you. This medicine helps to control your condition, but does not cure it.

It is important to keep taking your medicine even if you feel well.

## **DO NOT STOP TAKING TOPROL-XL TABLETS SUDDENLY.**

The dose needs to be reduced slowly over 7 to 14 days to make sure that your condition does not get worse. Your doctor will tell you how to gradually reduce the dose before stopping completely.

### **If you forget to take it**

**If you forget to take a dose of TOPROL-XL, skip the dose you missed and take your next dose when you are meant to.**

**Do not take a double dose to make up for the dose that you missed.**

**If you are not sure what to do, ask your doctor or pharmacist.**

If you have trouble remembering when to take your medicine, ask your pharmacist for some hints.

### **Overdose**

**Telephone your doctor, the Poisons Information Centre (13 11 26) or go to casualty at your nearest hospital immediately if you think that you or anyone else may have taken too much TOPROL-XL. Do this even if there are no**

**signs of discomfort or poisoning. You may need urgent medical attention.**

If you take too many TOPROL-XL tablets your blood pressure may drop too far. You will feel faint or may faint, and your heart rate will also slow down. You may also have nausea, vomiting and shortness of breath. In extreme cases, serious heart and lung problems may occur.

## **While you are using TOPROL-XL**

### **Things you must do**

**Be sure to keep all of your doctor's appointments so that your progress can be checked.**

**If you become pregnant while taking TOPROL-XL, tell your doctor immediately.**

**If you have a severe allergic reaction to foods, medicines or insect stings, tell your doctor immediately.**

If you have a history of allergies, there is a chance that TOPROL-XL may cause allergic reactions to be worse and harder to treat.

**If you feel light-headed, dizzy or faint when getting out of bed or standing up, get up slowly.**

You may feel light-headed or dizzy when you begin to take TOPROL-XL. This is because your blood pressure has fallen suddenly.

Standing up slowly, especially when you get up from bed or chairs, will help your body get used to the change in position and blood pressure. If this problem gets worse or continues, talk to your doctor.

**If you feel a worsening of your condition in the early stages of taking TOPROL-XL, tell your doctor immediately.**

Some people may experience an apparent worsening of their condition in the early stages of treatment with TOPROL-XL. It is important to tell your doctor if this happens to you, although it is usually temporary. If your condition continues to worsen, you should see your doctor as soon as possible.

**Make sure you drink enough water during exercise and hot weather when you are taking TOPROL-XL, especially if you sweat a lot.**

If you do not drink enough water while taking TOPROL-XL, you may feel faint or light-headed or sick. This is because your blood pressure is dropping too much. If you continue to feel unwell, tell your doctor.

**If you are being treated for diabetes, make sure you check your blood sugar level regularly and report any changes to your doctor.**

TOPROL-XL may change how well your diabetes is controlled. It may also cover up some of the symptoms of low blood sugar (hypoglycaemia). TOPROL-XL may increase the time your body takes to recover from low blood sugar. Your

doses of diabetic medicines, including insulin, may need to change.

**If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking TOPROL-XL.**

**Tell any doctors, dentists, and pharmacists who are treating you that you are taking TOPROL-XL.**

**If you plan to have surgery (even at the dentist) that needs an anaesthetic, tell your doctor or dentist that you are taking TOPROL-XL.**

**If you have to have any medical tests while you are taking TOPROL-XL, tell your doctor.**

TOPROL-XL may affect the results of some tests.

## **Things you must not do**

**Do not stop taking TOPROL-XL without checking with your doctor.**

Your doctor may want you to gradually reduce the amount of TOPROL-XL you are taking before stopping completely. This may help reduce the possibility of your condition worsening or other heart complications occurring.

**Do not give TOPROL-XL to anyone else even if they have the same condition as you.**

**Do not use TOPROL-XL to treat any other complaints unless your doctor tells you to.**

## **Things to be careful of**

**Be careful driving or operating machinery until you know how TOPROL-XL affects you.**

As with other beta-blocker medicines, TOPROL-XL may cause dizziness, light-headedness, tiredness, or drowsiness in some people. Make sure you know how you react to TOPROL-XL before you drive a car, operate machinery, or do anything else that could be dangerous if you are dizzy or light-headed.

**Be careful drinking alcohol while you are taking TOPROL-XL.**

If you drink alcohol, dizziness or light-headedness may be worse.

**Dress warmly during cold weather, especially if you will be outside for a long time (for example when playing winter sports).**

TOPROL-XL, like other beta-blocker medicines, tends to decrease blood circulation in the skin, fingers and toes. It may make you more sensitive to cold weather, especially if you have circulation problems.

## **Side effects**

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking TOPROL-XL.**

**If you get any side effects, do not stop taking TOPROL-XL without first talking to your doctor.**

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

If you are over 65 years of age you may have an increased chance of getting side effects.

**Ask your doctor or pharmacist to answer any questions you may have.**

**Tell your doctor if you notice any of the following and they worry you:**

- headache, tiredness, drowsiness, weakness, or lack of energy
- aches and pains, painful joints
- nausea (feeling sick), vomiting
- stomach upset, diarrhoea or constipation, weight gain
- dry mouth, changes in taste sensation
- difficulty sleeping, nightmares

- mood changes
- confusion, short-term memory loss, inability to concentrate
- increased sweating, runny or blocked nose
- hair loss

These side effects are usually mild.

**Tell your doctor or pharmacist immediately if you notice any of the following:**

- dizziness, light headedness or fainting especially on standing up, which may be a sign of low blood pressure
- tingling or "pins and needles"
- coldness, burning, numbness or pain in the arms and/or legs
- skin rash or worsening of psoriasis
- sunburn happening more quickly than usual
- abnormal thinking or hallucinations
- buzzing or ringing in the ears, deafness
- irritated eyes or blurred vision
- problems with sexual function

- constant "flu-like" symptoms with tiredness or lack of energy
- unusual bleeding or bruising.

These are serious side effects. You may need urgent medical attention. Serious side effects are rare.

**If any of the following happen, tell your doctor or go to Accident and Emergency at your nearest hospital immediately:**

- shortness of breath, being less able to exercise
- swelling of the ankles, feet or legs
- chest tightness, wheezing, noisy breathing, difficulty breathing
- chest pain, changes in heart rate or palpitations
- swelling of the face, lips, tongue or throat which may cause difficulty in swallowing or breathing, which may be signs of a serious allergic reaction
- yellowing of the skin or eyes (jaundice), generally feeling unwell.

These are very serious side effects. You may need urgent medical attention or hospitalisation. These side effects are rare.

**Other side effects not listed here may occur in some patients. Tell your doctor or pharmacist if you notice anything that is making you feel unwell.**

**Do not be alarmed by this list of possible side effects.**

You may not experience any of them.

## **After using TOPROL-XL**

### **Storage**

**Keep your tablets in the blister pack until it is time to take them.**

If you take TOPROL-XL out of the blister pack it will not keep well.

**Keep the tablets in a cool dry place where the temperature stays below 30°C.**

**Do not store it or any other medicine in the bathroom or near a sink. Do not leave it in the car or on window sills.**

Heat and dampness can destroy some medicines.

**Keep it where young children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

### **Disposal**

**Ask your pharmacist what to do with any tablets you have left over if your doctor tells you to stop taking them, or you find that the expiry date has passed.**

## Product description

### What TOPROL-XL looks like

TOPROL-XL 23.75 tablets are white to off-white, oval, scored on both sides and marked A/Beta Symbol on one side. This tablet can be divided into equal doses.

TOPROL-XL 47.5 tablets are white to off-white, round, scored on one side and marked A/mO on the other side. The score line is only there to help you break the tablet if you have difficulty swallowing it whole.

TOPROL-XL 95 tablets are white to off-white, round, scored on one side and marked A/mS on the other side. The score line is only there to help you break the tablet if you have difficulty swallowing it whole.

TOPROL-XL 190 tablets are white to off-white, oval, scored on one side and marked A/mY on the same side. The score line is only there to help you break the tablet if you have difficulty swallowing it whole.

### Ingredients

Each TOPROL-XL tablet contains metoprolol succinate 23.75mg, 47.5mg, 95mg or 190 mg as the active ingredient.

Plus the following inactive ingredients:

- Ethylcellulose

- Hyprollose
- Hypromellose
- Microcrystalline cellulose
- Synthetic paraffin
- Macrogol 6000
- Silicon dioxide
- Sodium stearyl fumarate
- Titanium dioxide

## **Manufacturer**

AstraZeneca Pty Ltd

ABN 54 009 682 311

66 Talavera Road

MACQUARIE PARK, NSW 2113

Telephone: 1800 805 342

## **Australian Registration Number:**

TOPROL-XL 23.75

Blister pack of 15 tablets

AUST R 97785

TOPROL-XL 47.5

Blister pack of 30 tablets

AUST R 97786

TOPROL-XL 95

Blister packs of 30 tablets

AUST R 97787

TOPROL-XL 190

Blister pack of 30 tablets

AUST R 97788

This leaflet was prepared in October 2017.

TOPROL-XL is a trade mark of the AstraZeneca group of companies

© AstraZeneca 2017

Doc ID-001272100 V3.0